


CLOBETASOL PROPIONATE ANDA- clobetasol propionate ointment **Macleods Pharmaceuticals Limited**

Clobetasol Propionate Ointment USP, 0.05% **FOR TOPICAL DERMATOLOGIC USE ONLY NOT FOR OPHTHALMIC, ORAL, OR** **INTRAVAGINAL USE**

DESCRIPTION

Clobetasol propionate ointment USP, contain the active compound clobetasol propionate, a synthetic corticosteroid, for topical dermatologic use. Clobetasol, an analog of prednisolone, has a high degree of glucocorticoid activity and a slight degree of mineralocorticoid activity.

Chemically, clobetasol propionate is (11 β , 16 β)-21-chloro-9-fluoro-11-hydroxy-16-methyl-17-(1-oxopropoxy)-pregna-1,4-diene-3,20-dione, and it has the following structural formula:



Clobetasol propionate has the empirical formula C₂₅H₃₂ClF₀₅ and a molecular weight of 467. It is a white to cream-colored crystalline powder insoluble in water.

Clobetasol propionate ointment USP, clobetasol propionate 0.5 mg/g in a base of propylene glycol, sorbitan sesquioleate, and white petrolatum.

CLINICAL PHARMACOLOGY

Like other topical corticosteroids, clobetasol propionate has anti-inflammatory, antipruritic, and vasoconstrictive properties. The mechanism of the anti-inflammatory activity of the topical steroids, in general, is unclear.

However, corticosteroids are thought to act by the induction of phospholipase A₂ inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor, arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A₂.

Pharmacokinetics

The extent of percutaneous absorption of topical corticosteroids is determined by many factors, including the vehicle and the integrity of the epidermal barrier. Occlusive dressing with hydrocortisone for up to 24 hours has not been demonstrated to increase penetration; however, occlusion of hydrocortisone for 96 hours markedly enhances penetration.

Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin may increase percutaneous absorption.

Studies performed with clobetasol propionate ointment indicate that they are in the super-high range of potency as compared with other topical corticosteroids.

INDICATIONS & USAGE

Clobetasol propionate ointment is super-high potency corticosteroid formulations indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses. Treatment beyond 2 consecutive weeks is not recommended, and the total dosage should not exceed 50 g/week because of the potential for the drug to suppress the hypothalamic-pituitary-adrenal (HPA) axis. Use in pediatric patients under 12 years of age is not recommended.

As with other highly active corticosteroids, therapy should be discontinued when control has been achieved. If no improvement is seen within 2 weeks, reassessment of the diagnosis may be necessary.

CONTRAINDICATIONS

Clobetasol propionate ointment is contraindicated in those patients with a history of hypersensitivity to any of the components of the preparations.

PRECAUTIONS

General:

Clobetasol propionate ointment should not be used in the treatment of rosacea or perioral dermatitis, and should not be used on the face, groin, or axillae.

Systemic absorption of topical corticosteroids can produce reversible HPA axis suppression with the potential for glucocorticosteroid insufficiency after withdrawal from treatment.

Manifestations of Cushing's syndrome, hyperglycemia, and glucosuria can also be produced in some patients by systemic absorption of topical corticosteroids while on therapy.

Patients applying a topical steroid to a large surface area or to areas under occlusion should be evaluated periodically for evidence of HPA axis suppression. This may be done by using the ACTH stimulation, A.M. plasma cortisol, and urinary free cortisol tests.

Patients receiving super-potent corticosteroids should not be treated for more than 2 weeks at a time, and only small areas should be treated at any one time due to the increased risk of HPA suppression.

Clobetasol propionate ointment produced HPA axis suppression when used at doses as low as 2 g/day for 1 week in patients with eczema.

If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent corticosteroid.

Recovery of HPA axis function is generally prompt upon discontinuation of topical corticosteroids. Infrequently, signs and symptoms of glucocorticosteroid insufficiency may occur that require supplemental systemic corticosteroids. For information on systemic supplementation, see prescribing information for those products.

Pediatric patients may be more susceptible to systemic toxicity from equivalent doses due to their larger skin surface to body mass ratios (see PRECAUTIONS: Pediatric Use). If irritation develops, clobetasol propionate ointment should be discontinued and appropriate therapy instituted. Allergic contact dermatitis with corticosteroids is usually diagnosed by observing a failure to heal rather than noting a clinical exacerbation as with most topical products not containing corticosteroids. Such an observation should be corroborated with appropriate diagnostic patch testing.

If concomitant skin infections are present or develop, an appropriate antifungal or antibacterial agent should be used.

If a favorable response does not occur promptly, use of clobetasol propionate ointment should be discontinued until the infection has been adequately controlled.

Information for Patients:

Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
2. This medication should not be used for any disorder other than that for which it was prescribed.
3. The treated skin area should not be bandaged, otherwise covered, or wrapped so as to be occlusive unless directed by the physician.
4. Patients should report any signs of local adverse reactions to the physician.

Laboratory Tests:

The following tests may be helpful in evaluating patients for HPA axis suppression:

ACTH stimulation test

A.M. plasma cortisol test

Urinary free cortisol test

Carcinogenesis, Mutagenesis, Impairment of Fertility:

Long-term animal studies have not been performed to evaluate the carcinogenic potential of clobetasol propionate.

Studies in the rat following subcutaneous administration at dosage levels up to 50 mcg/kg per day revealed that the females exhibited an increase in the number of resorbed embryos and a decrease in the number of living fetuses at the highest dose. Clobetasol propionate was nonmutagenic in 3 different test systems: the Ames test, the *Saccharomyces cerevisiae* gene conversion assay, and the *E coli* B WP2 fluctuation test.

Pregnancy: Teratogenic Effects:

Corticosteroids have been shown to be teratogenic in laboratory animals when administered systemically at relatively low dosage levels. Some corticosteroids have been shown to be teratogenic after dermal application to laboratory animals.

Clobetasol propionate has not been tested for teratogenicity when applied topically; however, it is absorbed percutaneously, and when administered subcutaneously it was a significant teratogen in both the rabbit and mouse. Clobetasol propionate has greater teratogenic potential than steroids that are less potent.

Teratogenicity studies in mice using the subcutaneous route resulted in fetotoxicity at the highest dose tested (1 mg/kg) and teratogenicity at all dose levels tested down to 0.03 mg/kg. These doses are approximately 1.4 and 0.04 times, respectively, the human topical dose of clobetasol propionate ointment. Abnormalities seen included cleft palate and skeletal abnormalities.

In rabbits, clobetasol propionate was teratogenic at doses of 3 and 10 mcg/kg. These doses are approximately 0.02 and 0.05 times, respectively, the human topical dose of clobetasol propionate ointment. Abnormalities seen included cleft palate, cranioschisis, and other skeletal abnormalities.

There are no adequate and well-controlled studies of the teratogenic potential of clobetasol propionate in pregnant women. Clobetasol propionate ointment should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers:

Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in human milk. Because many drugs are excreted in human milk, caution should be exercised when clobetasol propionate ointment is administered to a nursing woman.

Pediatric Use:

Safety and effectiveness of clobetasol propionate ointment in pediatric patients have not been established. Use in pediatric patients under 12 years of age is not recommended. Because of a higher ratio of skin surface area to body mass, pediatric patients are at a greater risk than adults of HPA axis suppression and Cushing's syndrome when they are treated with topical corticosteroids. They are therefore also at greater risk of adrenal insufficiency during or after withdrawal of treatment. Adverse effects including striae have been reported with inappropriate use of topical corticosteroids in infants and children.

HPA axis suppression, Cushing's syndrome, linear growth retardation, delayed weight gain, and intracranial hypertension have been reported in children receiving topical corticosteroids.

Manifestations of adrenal suppression in children include low plasma cortisol levels and an absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Geriatric Use:

A limited number of patients at or above 65 years of age have been treated with clobetasol propionate ointment (n = 101) in US and non-US clinical trials. While the number of patients is too small to permit separate analysis of efficacy and safety, the adverse reactions reported in this population were similar to those reported by younger patients. Based on available data, no adjustment of dosage of clobetasol propionate ointment in geriatric patients is warranted.

ADVERSE REACTIONS

In controlled clinical trials, the most frequent adverse events reported for clobetasol propionate ointment were burning sensation, irritation, and itching in 0.5% of treated patients. Less frequent adverse reactions were stinging, cracking, erythema, folliculitis, numbness of fingers, skin atrophy, and telangiectasia.

Cushing's syndrome has been reported in infants and adults as a result of prolonged use of topical clobetasol propionate formulations.

The following additional local adverse reactions have been reported with topical corticosteroids, and they may occur more frequently with the use of occlusive dressings and higher potency corticosteroids. These reactions are listed in an approximately decreasing order of occurrence: dryness, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, secondary infection, irritation, striae, and miliaria.

OVERDOSAGE:

Topically applied clobetasol propionate ointment can be absorbed in sufficient amounts to produce systemic effects (see PRECAUTIONS).

DOSAGE & ADMINISTRATION:

Apply a thin layer of clobetasol propionate ointment USP, to the affected skin areas twice daily and rub in gently and completely (see INDICATIONS AND USAGE).

Clobetasol propionate ointment USP, is super-high potency topical corticosteroids; therefore, treatment should be limited to 2 consecutive weeks and amounts greater than 50 g/week should not be used.

As with other highly active corticosteroids, therapy should be discontinued when control has been achieved. If no improvement is seen within 2 weeks, reassessment of diagnosis may be necessary.

Clobetasol propionate ointment USP, should not be used with occlusive dressings.

Geriatric Use:

In studies where geriatric patients (65 years of age or older, see PRECAUTIONS) have been treated with clobetasol propionate ointment safety did not differ from that in younger patients; therefore, no dosage adjustment is recommended.

HOW SUPPLIED:

Clobetasol propionate ointment USP, 0.05% is supplied in

15 g tube NDC 33342-506-15

30 g tube NDC 33342-506-30

45 g tube NDC 33342-506-45

60 g tube NDC 33342-506-60

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

Do not refrigerate.


Manufactured for :
Macleods Pharma USA, Inc.
Princeton, NJ 08540

Manufactured by:
Macleods Pharmaceuticals Ltd.
At Oxalis Labs, Baddi, Himachal Pradesh, INDIA

Revised: January 2024

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL


Clobetasol Propionate Ointment USP, 0.05% w/w
Pack Count: 15 g Tube
NDC 33342-506-15




MACLEODS
Pharma USA
NDC 33342-506-15
Clobetasol Propionate Ointment USP,
0.05 %
FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE. NET WT 15 grams
Rx Only

<p>Each gram contains: Clobetasol propionate, USP 0.5 mg in a base of propylene glycol, sorbitan sesquioleate, and white petrolatum.</p> <p>USUAL DOSAGE: See package insert for full prescribing information.</p> <p>WARNING: Keep out of reach of children.</p> <p>TO OPEN: Use cap to puncture seal.</p> <p>IMPORTANT: Do not use if seal has been punctured or is not visible.</p>	<p>Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].</p> <p>DO NOT REFRIGERATE.</p> <p>Code No.: HP/DRUGS/12/665</p> <p>MACLEODS Manufactured for : Macleods Pharma USA, Inc. Princeton, NJ 08540</p>	<p>PM00373101</p> <p>See LOT and EXP. ▸</p>  <p>Manufacturer : Macleods Pharmaceuticals Ltd. At Oxalis Labs Baddi, Himachal Pradesh, INDIA</p>
---	---	--

Clobetasol Propionate Ointment USP, 0.05% w/w
Pack Count: 15 g Carton
NDC 33342-506-15



Clobetasol Propionate Ointment USP, 0.05% w/w
 Pack Count: 30 g Tube
 NDC 33342-506-30



NDC 33342-506-30

Clobetasol Propionate Ointment USP,

0.05 %

Rx Only

FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE.

NET WT 30 grams

Each gram contains:

Clobetasol propionate, USP 0.5 mg in a base of propylene glycol, sorbitan sesquioleate, and white petrolatum.

USUAL DOSAGE: See package insert for full prescribing information.

WARNING:

Keep out of reach of children.

TO OPEN: Use cap to puncture seal.

IMPORTANT:

Do not use if seal has been punctured or is not visible.

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]. **DO NOT REFRIGERATE.**

Code No.:

HP/DRUGS/12/665



Manufactured for :

Macleods Pharma USA, Inc.
Princeton, NJ 08540

Manufacturer :


Macleods Pharmaceuticals Ltd.
At Oxalis Labs
Baddi, Himachal Pradesh, INDIA

See LOT and EXP. Date ↗




PM00373301

Clobetasol Propionate Ointment USP, 0.05% w/w
Pack Count: 30 g Carton
NDC 33342-506-30



Clobetasol Propionate Ointment USP, 0.05% w/w
Pack Count: 45 g Tube
NDC 33342-506-45



MACLEOD'S
Pharma USA
NDC 33342-506-45

Clobetasol Propionate Ointment USP,

0.05 %

FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE.

NET WT 45 grams

R_x Only

Each gram contains:
Clobetasol propionate, USP 0.5 mg
in a base of propylene glycol, sorbitan
sesquioleate, and white petrolatum.

USUAL DOSAGE: See package
insert for full prescribing information.

WARNING:
Keep out of reach of children.

TO OPEN: Use cap to puncture seal.


IMPORTANT:
Do not use if seal has been
punctured or is not visible.

Store at 20° to 25°C (68° to 77°F);
excursions permitted to 15° to 30°C
(59° to 86°F) [See USP Controlled
Room Temperature].
DO NOT REFRIGERATE.

MACLEOD'S
Manufactured for :
Macleods Pharma USA, Inc.
Princeton, NJ 08540

Manufacturer :
Macleods Pharmaceuticals Ltd.
At Oxalis Labs
Baddi, Himachal Pradesh, INDIA

**See LOT and
EXP. Date** ➔




Code No.:
HP/DRUGS/12/665
PM00373501

Clobetasol Propionate Ointment USP, 0.05% w/w
Pack Count: 45 g Carton
NDC 33342-506-45



Clobetazol Propionate Ointment USP, 0.05% w/w
 Pack Count: 60 g Tube
 NDC 33342-506-60



MACLEODS
Pharma USA

NDC 33342-506-60

Clobetasol Propionate Ointment USP,

0.05 %

FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE.

R_x Only

NET WT 60 grams

Each gram contains:
Clobetasol propionate, USP 0.5 mg
in a base of propylene glycol, sorbitan
sesquioleate, and white petrolatum.

USUAL DOSAGE: See package
insert for full prescribing information.

WARNING:
Keep out of reach of children.

TO OPEN: Use cap to puncture seal.

Store at 20° to 25°C (68° to 77°F);
excursions permitted to 15° to 30°C
(59° to 86°F) [See USP Controlled
Room Temperature].

DO NOT REFRIGERATE.

IMPORTANT: Do not use if seal has
been punctured or is not visible.

MACLEODS

Manufactured for :
Macleods Pharma USA, Inc.
Princeton, NJ 08540


Code No.: HP/DRUGS/12/665
PM00373701

See LOT and EXP. Date →



Manufacturer :
Macleods Pharmaceuticals Ltd.
At Oxalis Labs
Baddi, Himachal Pradesh, INDIA

Clobetasol Propionate Ointment USP, 0.05% w/w
Pack Count: 60 g Carton
NDC 33342-506-60



CLOBETASOL PROPIONATE ANDA

clobetasol propionate ointment

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:33342-506
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CLOBETASOL PROPIONATE (UNII: 779619577M) (CLOBETASOL - UNII:ADN79D536H)	CLOBETASOL PROPIONATE	0.5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITAN SESQUIOLEATE (UNII: 0W8RRI5W5A)	
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date

1	NDC:33342-506-15	1 in 1 CARTON	12/28/2023	
1		15 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:33342-506-30	1 in 1 CARTON	12/28/2023	
2		30 g in 1 TUBE; Type 0: Not a Combination Product		
3	NDC:33342-506-45	1 in 1 CARTON	12/28/2023	
3		45 g in 1 TUBE; Type 0: Not a Combination Product		
4	NDC:33342-506-60	1 in 1 CARTON	12/28/2023	
4		60 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215990	12/28/2023	

Labeler - Macleods Pharmaceuticals Limited (862128535)

Establishment

Name	Address	ID/FEI	Business Operations
OXALIS LABS		860120472	ANALYSIS(33342-506) , LABEL(33342-506) , MANUFACTURE(33342-506) , PACK(33342-506)

Revised: 1/2024

Macleods Pharmaceuticals Limited