

FAMILY WELLNESS PAIN RELIEF- benzocaine 20% gel
FAMILY DOLLAR (Family Wellness)

Family Wellness Oral Pain Relief

Drug Fact

Active Ingredient

Benzocaine 20%

Purpose

Oral Pain Reliever

Uses

- temporary relief of occasional minor irritation, pain and sore mouth

Warnings

Methemoglobinemia Warning: Use of this product may cause Methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops;

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

more than directed

for more than 7 days unless told to do so by a dentist or doctor

Stop use and ask a Doctor if

swelling, rash or fever develops

irritation, pain, or redness persists or worsens

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not use tube if it is cut prior to opening
- cut open tip of tube on score mark
- use your fingertip or cotton applicator to apply a small pea-size amount of Oral Pain Relief Gel
- apply to affected area up to four times daily or as directed by a dentist or physician
- Adults and children 2 years of age and older: Apply to affected area
- Children under 12 years of age should be supervised in the use of this product
- Children under 2 years of age: Consult a doctor

Inactive Ingredients

Benzyl Alcohol, Carbomer, D&C Yellow No 10, FD&C Blue No 1, FD&C Red No 40, Flavor, Glycerin, Methylparaben, Polyethylene Glycol, Propylene Glycol, Sodium Saccharin

Other information

store at a controlled room temperature 68°- 77°F (20°-25°C)

Principal Display Panel

NDC 55319-157-33

Family Wellness

Oral Pain Relief

MAXIMUM STRENGTH GEL

Benzocaine 20% NET WT. 0.33 OZ (9.35g)

FAMILY
Wellness

COMPARE TO THE
ACTIVE INGREDIENT IN
MAXIMUM STRENGTH
ANBESOL® GEL*

MAXIMUM STRENGTH

ORAL ANESTHETIC GEL

BENZOCAINE 20%

Oral pain relief gel

NET WT 0.33 OZ (9.35 g)

NDC 55319-157-33

FAST-ACTING PAIN RELIEF FROM:

- Toothaches
- Gum pain
- Canker sores
- Dentures

FAMILY
Wellness

COMPARE TO THE
ACTIVE INGREDIENT IN
MAXIMUM STRENGTH
ANBESOL® GEL*

MAXIMUM STRENGTH

ORAL ANESTHETIC GEL

BENZOCAINE 20%

Oral pain relief gel

NET WT 0.33 OZ (9.35 g)

NDC 55319-157-33



MADE IN THE U.S.A. OF U.S. AND IMPORTED
INGREDIENTS AND COMPONENTS
510 VOLVO PKWY., CHESTERPAKE, VA 23320 USA
DISTRIBUTED BY: MIDWOOD BRANDS, LLC
NOT 100% SATISFIED?
Return with in 30 days to the
store of purchase for a refund
(with receipt) or exchange.
This product is not
distributed by
Consumer Brands
Foundation
over of the
registered trademark
Anbesol®.

<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash or fever develops <p>Keep out of the reach of children.</p> <p>Directions</p> <ul style="list-style-type: none"> • adults and children 2 years of age or older: apply to affected area up to 4 times daily or as directed by doctor/dentist • children between 2 and 12 years of age: adult supervision should be given in use of this product • children under 2 years of age: do not use <p>Other information</p> <ul style="list-style-type: none"> • do not use tube if it is cut prior to opening • to open tube, cut open tip of tube on score mark with scissors • store at a controlled room temperature 68°-77°F (20°-25°C) <p>Inactive ingredients</p> <p>Benzyl Alcohol, Carbomer, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, Flavor, Glycerin, Methylparaben, Polyethylene Glycol, Propylene Glycol, Sodium Saccharin</p> <p>Questions? Call 800-814-8028</p>	<p>Drug Facts</p> <p>Active ingredient</p> <p>Benzocaine 20% Oral anesthetic</p> <p>Purpose</p> <p>For the temporary relief of pain associated with:</p> <ul style="list-style-type: none"> • toothache • canker sores • minor dental procedures • minor injury of the mouth and gums • caused by dentures or orthodontic appliances <p>Warnings METHEMOGLOBINEMIA WARNING: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:</p> <ul style="list-style-type: none"> • pale, gray, or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy <p>Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics. If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Do not use</p> <ul style="list-style-type: none"> • for teething • in children under 2 years of age <p>When using this product</p> <ul style="list-style-type: none"> • avoid contact with the eyes • do not exceed recommended dosage • do not use for more than 7 days unless told to do so by a dentist or doctor
---	---

FAMILY
Wellness
MAXIMUM STRENGTH
ORAL ANESTHETIC

557131



COMPARE TO THE ACTIVE INGREDIENT IN
MAXIMUM STRENGTH ANBESOL® GEL*

MAXIMUM STRENGTH

ORAL ANESTHETIC

GEL

BENZOCAINE 20%

Oral pain relief gel

NET WT 0.33 OZ (9.35 g)

NDC 55319-157-33

Active ingredient Benzocaine 20% **Purpose** Oral anesthetic **Uses** for the temporary relief of pain associated with:
• toothache • canker sores • minor dental procedures • minor injury of the mouth and gums caused by dentures or
orthodontic appliances **Warnings** **METHEMOGLOBINEMIA WARNING:** Use of this product may cause
methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in
blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a
child in your care develops: • pale, gray, or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath
• dizziness or lightheadedness • fatigue or lack of energy **Allergy alert:** do not use this product if you have a history of
allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics. If a skin reaction occurs,
stop use and seek medical help right away. **Do not use** • for teething • in children under 2 years of age **When using this
product** • avoid contact with the eyes • do not exceed recommended dosage • do not use for more than 7 days unless
told to do so by a dentist or doctor **Stop use and ask a doctor if** • sore mouth symptoms do not improve in 7 days
• irritation, pain, or redness persists or worsens • swelling, rash or fever develops **Keep out of the reach of children.** In
case of overdose, get medical help or contact a Poison Control Center right away. **Directions** • adults and children 2 years
of age or older: apply to affected area up to 4 times daily or as directed by doctor/dentist • children between 2 and 12 years
of age: adult supervision should be given in use of this product • children under 2 years of age: do not use **Other
information** • do not use tube if it is cut prior to opening • to open tube, cut open tip of tube on score mark with scissors
• store at a controlled room temperature 68°-77°F (20°-25°C) **Questions? Call** 800-814-8028
DISTRIBUTED BY: MIDWOOD BRANDS, LLC, 510 VOLVO PKWY., CHESAPEAKE, VA 23320 USA
* This product is not manufactured or distributed by Foundation Consumer Brands,
owner of the registered trademark Anbesol®.

FAMILY WELLNESS PAIN RELIEF

benzocaine 20% gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-157
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
CARBOMER 940 (UNII: 4Q93RCW27E)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN (UNII: A2I8C7HI9T)	

POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
SODIUM SACCHARIN (UNII: SB8ZUX40TY)				
Product Characteristics				
Color		Score		
Shape		Size		
Flavor	MINT	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-157-33	1 in 1 CARTON	01/15/2026	
1		9.35 g in 1 TUBE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022		01/15/2026	

Labeler - FAMILY DOLLAR (Family Wellness) (024472631)

Registrant - Sheffield Pharmaceuticals LLC (151177797)

Establishment

Name	Address	ID/FEI	Business Operations
Sheffield Pharmaceuticals LLC		151177797	manufacture(55319-157) , analysis(55319-157)

Revised: 1/2026

FAMILY DOLLAR (Family Wellness)