

7.5% HYPERTONIC SALINE- hypertonic saline solution liquid

Radix Laboratories, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

7.5% Hypertonic Saline

• CONTENTS (grams/100ml):

- Sodium Chloride7.5

Inactives:

- Purified Water q.s.
- Methylparaben
- Propylparaben
- Citric Acid

Indications:

- For use as a supplemental nutritive source of sodium chloride in cattle.

Storage:

- Store at controlled room temperature between 59°F to 86°F (15°C -30°C).

Warnings:

- Avoid storage at cold temperatures which can cause product to become cloudy.
- To clarify place product in warm water bath (100°F-120°F) for about 30 minutes.

Dosage and administration:

- Administer orally as a drench or by use of a stomach tube.
- The usual recommended dose in adult cattle is 16.9 fl. oz (500mL) to 33.8 fl.oz (1000mL) depending on size and condition.
- Keep away from children.

PRINCIPAL DISPLAY PANEL

CONTENTS (grams/100mL):

Sodium Chloride 7.5
Purified Water q.s.
Inactives: Methylparaben, propylparaben,
Citric Acid

STORAGE:

Store at controlled room temperature
between 59°F to 86°F (15°C to 30°C)

MANUFACTURED BY:

Radix Laboratories, Inc.
1334 International Drive
Eau Claire, WI 54701
(800) 927-2349

TAKE TIME  READ LABEL
INSTRUCTIONS



NDC 50480-124-99



**7.5%
HYPERTONIC
SALINE**

FOR ANIMAL USE ONLY

**NET CONTENTS
33.8 FL. OZ. (1000 ML)**

WARNING:

Avoid prolonged storage at cold temperatures,
which can cause preservatives to precipitate.
To clarify place product in warm water bath
(100°F -120°F) for about 30 minutes.

INDICATIONS:

For use as a supplemental nutritive source of
sodium chloride in cattle.

Dosage and Administration:

Administer orally as a drench. The usual
recommended dose in adult cattle is 500 to
1000 ml, depending on size and condition.

Keep away from children.

Item image



NDC 50480-124-99

Radix
Labs

**7.5%
HYPERTONIC
SALINE**

FOR ANIMAL USE ONLY

**NET CONTENTS
33.8 FL. OZ. (1000 ML)**

WARNING:

Avoid prolonged storage at room temperature which can cause precipitation. To clarify place product in water bath (100°F - 120°F) for about 30 minutes.

INDICATIONS:

For use as a supplement to sodium chloride in cattle.

Dosage and Administration:

Administer orally as a bolus. Recommended dose is 1000 ml, depending on animal weight.

Keep away from children.

Contents (100mL):

7.5

q.s.

propylparaben,

temperature

(100°F to 30°C)

BY:

, inc.

Drive

48701

READ LABEL

INSTRUCTIONS



LOT NO. R3E105
EXP. DATE 05/2018

7.5% HYPERTONIC SALINE

hypertonic saline solution liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:50480-124
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X) (CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	7.5 g in 100 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50480-124-99	12 in 1 CARTON		
1		1000 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/27/2013	

Labeler - Radix Laboratories, Inc. (108500091)

Revised: 12/2014

Radix Laboratories, Inc.