RANITIDINE- ranitidine hydrochloride tablet, film coated Supervalu Inc.

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Ranitidine 75 mg (as ranitidine hydrochloride, USP 84 mg)

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

OTHER INFORMATION

- TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.
- store at 20° 25° C (68° 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

INACTIVE INGREDIENTS


Colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, talc, titanium dioxide

QUESTIONS?

Call **1-877-932-7948**

PRINCIPAL DISPLAY PANEL NDC 41163-353-31 EQUALINE Compare to Zantac 75[®]Active Ingredient^{*} Acid Reducer ranitidine Tablets, USP 75 mg (acid reducer) prevents & relieves heartburn associated with acid indigestion & sour stomach 30 tablets DISTRIBUTED BY: SUPERVALU INC.

5115091 03/15



HYPROMELLOSES (UNII: 3NXW29V3WO)							
FERRIC OXIDE RED (UNII: 1K09F3G675)							
MAGNESIUM STEARATE (UNII: 70097M6I30)							
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)							
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)							
TALC (UNII: 7SEV7J4R1U)							
TITANIUM DIO XIDE	UNII: 15F	FIX9 V2JP)					
Product Characte	eristics						
Color	Color		Score		no score		
Shape	Shape		Size		8 mm		
Flavor			Imprint Code		OR;606		
Contains							
Packaging							
Packaging # Item Code		Package Descrip	otion	Marketing Start Date	Marketing End Date		
0.0	30 in 1 B	Package Descrip OTTLE; Type 0: Not a Col		Marketing Start Date	Marketing End Date		
# Item Code	30 in 1 E			Marketing Start Date	• Marketing End Date		
# Item Code	30 in 1 E			Marketing Start Date	Marketing End Date		
# Item Code		SOTTLE; Type 0: Not a Con		Marketing Start Date	Marketing End Date		
# Item Code 1 NDC:41163-353-31	orma	SOTTLE; Type 0: Not a Con	mbination Product	Marketing Start Date			
 # Item Code 1 NDC:41163-353-31 Marketing Inf 	orma y Apj	SOTTLE; Type 0: Not a Con	mbination Product				

Labeler - Supervalu Inc. (006961411)

Registrant - Ranbaxy Pharmaceuticals Inc. (184769029)

Establishment

Name	Address	ID/FEI	Business Operations
Shasun Pharmaceuticals Limited		915786829	MANUFACTURE(41163-353)

Revised: 4/2015

Supervalu Inc.