## CAREALL PAIN RELIEF PM EXTRA STRENGTH- acetaminophen, diphenhydramine hcl tablet New World Imports, Inc

\_\_\_\_\_

## **Drug Facts**

## **Active Ingredients**

Acetaminophen 500mg Diphenhydramine HCl 25mg

## Purpose

Pain reliever Nightime sleep aid

## Keep Out of reach of Children

**Keep out of reach of children.** In case of accidental overdose, contact a doctor or Poison Control Center (1-800-222-1222) right away. Quick Medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Uses

Temporarily relieves occasional headaches and minor aches and pains with accompanying sleeplessness

## Warnings

**Liver Warning**: This product contains Acetaminophen. Severe liver damage may occur if you take

- More than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

**Allergy Alert:**Acetaminophen may cause sever skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

## Do not use:

- in children under 12 years of age
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- if you have even had an allergic reaction to this product or any of its ingredients

## Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

## Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

# When using this product

- drowsiness will occur
- •do not drive a motor vehicle or operate machinery
- avoid alcoholic drinks

# Stop use and ask a doctor if

•sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness

- •pain gets worse or last for more than 10 days.
- •fever gets worse or last for more than 3 days
- •redness or swelling are present

•new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

## Directions


# Do not use more than directed. (see overdose warning)

Adults and children 12 years of age and older: Take 2 caplets at bedtime. Do not take more than 2 caplets in 24 hours

Children under 12 years: do not use

# Inactive Ingredients

colloidal silicon dioxide, copovidone, croscarmellose sodium, FD and C blue 1, hypromellose, lactose, magnesium, stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide



S	Questions or comments? Call 1-844-705-4384 Mon - Fri	Inactive ingredi linactive, copovidone, blue #1, hypromellos stearate, microcryst glycol, povidone, pre glycol, povidone, stea starch glycolate, stea	• Store between 20-25°C	Directions do not take more that adults and childred 2 caplets at bedti 2 caplets of this j children under 12	If pregnant or breast-f professional before use, children. Overdose warn get medical help or conta get medical help or conta right away (1-800-222- right away (1-800-222- attention is critical for add even if you do not notice		<ul> <li>Stop use and ask a doctor if sleeplessness persists continue than 2 weeks, insomnia ma of serious underlying medic prior det worce or loster mo prior det worce or loster mo</li> </ul>	<ul> <li>When using this product</li> <li>drowsiness will occur</li> <li>do not drive a motor vehic</li> </ul>	Ask a doctor or pharmacist before taking the blood thinning drug w taking sedatives or tranquilizers	<ul> <li>Ask a doctor before use if you have</li> <li>a breathing problem such as er chronic bronchitis</li> <li>travia to prochere</li> </ul>	Drug Facts (continued)
STOP PEELING	i 8:30 AM -	<i>Inactive ingredients</i> colloidal silicon dioxide, copovidone, croscarmellose sodium, FD&C blue #1, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide	<b>-25°C (68-77°F) in a dry</b>	Directions • do not take more than directed (see overdose warning) • adults and children 12 years and over: take 2 caplets at bedtime; do not take more than 2 caplets of this product in 24 hours • children under 12 years: do not use	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	<ul> <li>fever gets worse or lasts more than 3 days</li> <li>fever gets worse or lasts more than 3 days</li> <li>redness or swelling is present</li> <li>new symptoms occur</li> <li>These could be signs of a serious condition.</li> </ul>	op use and ask a doctor if sleeplessness persists continuously for more than 2 weeks. Insommia may be a symptom of serious underlying medical illness.	Intern using this product drowsiness will occur avoid alcoholic drinks do not drive a motor vehicle or operate machinery	todute unitating due to an enlarged prostate giant Ask a doctor or pharmacist before use if you are     taking the blood thinning drug warfarin     taking sedatives or tranquilizers	n ph =	ntinued)
	5 PM EST	con m, FD&C n nylene dium dium	dry place	e warning) : take e than	ealth ach of overdose, ol Center idical r children mptoms.	days ion.	nptom	lic drinks nachinery	n n n n n n n n n n n n n n n n n n n	liver disease ysema or glaucoma	

# CAREALL PAIN RELIEF PM EXTRA STRENGTH

acetaminophen, diphenhydramine hcl tablet

. . . .

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51824-108
Route of Administration	ORAL		

Active Ingredient/Active Moiety						
Ingredient Name	<b>Basis of Strength</b>	Strength				
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	500 mg				
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg				

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

		_					
POVIDONE (UNII: FZ		-					
STARCH, CORN (UN		-					
STEARIC ACID (UNII							
TITANIUM DIOXIDE		-					
SILICON DIOXIDE (	-						
			(UNII: AG9B65PV6B)				
COPOVIDONE K25-							
LACTOSE, UNSPEC	IFIED FOR	<b>M</b> (UNII: J2B2A4	N98G)				
MAGNESIUM STEAF	•	•					
POLYETHYLENE GL	YCOL, UN	SPECIFIED (UN	II: 3WJQ0SDW1A)				
Product Chara	cteristic	CS					
Color		blue	Score	no score			
Shape		OVAL	Size	17mm			
Flavor			Imprint Code	AAA;1031			
Contains							
Packaging							
		Package Description		Marketing Start	Ma	Marketing End Date	
# Item Code		Package Des	scription	Date			
NDC:51824_108		-	•	-		-	
NDC:51824-108-	50 in 1 BO	-	•	Date			
<b>1</b> NDC:51824-108- 50	50 in 1 BO Product	TTLE; Type 0: N	•	Date			
<b>1</b> NDC:51824-108- 50	50 in 1 BO Product	TTLE; Type 0: N	•	Date			
NDC:51824-108-	50 in 1 BO Product <b>nform</b>	TTLE; Type 0: N	ot a Combination er or Monograph	<b>Date</b> 08/23/2024		_	

Labeler - New World Imports, Inc (07537227	6)
--	----

Revised: 8/2024

New World Imports, Inc