

LOHIST D- chlorpheniramine maleate / pseudoephedrine hcl liquid
Larken Laboratories, Inc.

LoHist D

Drug Facts

Active Ingredients

(In each 5 mL teaspoonful)

Chlorpheniramine Maleate, USP 2 mg

Pseudoephedrine HCl, USP 30 mg

Purpose

Chlorpheniramine Maleate Antihistamine

Pseudoephedrine HCl Nasal decongestant

Uses

temporarily relieves these symptoms due to hay fever (allergic rhinitis):

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies.
- temporarily restores freer breathing through the nose

Warnings

Do not use

- to sedate a child or make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this drug.

Ask a Doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes

- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are.

- taking any other nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product

Do not exceed recommended dosage.

- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur.
- symptoms do not improve within 7 days or are accompanied by fever.

If pregnant or breast feeding

ask a health professional before use.

Keep out of the reach of children

In case of overdose, get medical help or contact a Poison Control Center immediately.

Directions

Do not exceed 4 doses in a 24-hour period

Age	Dose
Adults and children over 12 years of age	2 teaspoonfuls (10 mL) every 4 to 6 hours
Children 6 to under 12 years of age	1 teaspoonful (5 mL) every 4 to 6 hours
Children under 6 years of age	Ask your doctor

Other information

- store at 20°-25°C (68°-77°F)
- very low sodium, contains less than 1 mg sodium per 5 mL teaspoonful

Inactive ingredients

cherry flavoring, methylparaben, polyethylene glycol, propylparaben, purified water saccharin sodium, and sorbitol

Questions or comments?

Call 1-601-855-7678 weekdays from 9:00 am to 4:00 pm CST or go to <http://www.larkenlabs.com>.

Package Label, Principal Display Panel

Figure 1:16 oz. bottle label

NDC 68047-120-16

LoHIST-D

ANTIHISTAMINE
NASAL DECONGESTANT

SUGAR FREE / ALCOHOL FREE
DYE FREE

Cherry Flavored Liquid

**DO NOT USE IF FOIL SEAL UNDER
THE CAP IS BROKEN OR MISSING.**

Distributed by:
**LARKEN
LABORATORIES**
Canton, MS 39046

16 fl. oz. (473 mL)



Lot/Exp. date:

Drug Facts

Active ingredients	Purpose
<i>(in each 5 mL teaspoonful)</i> Chlorpheniramine Maleate, USP 2 mg Antihistamine Pseudoephedrine HCl, USP 30 mg ... Nasal decongestant	

Uses
temporarily relieves these symptoms due to hay fever (allergic rhinitis): ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes ■ temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies ■ temporarily restores freer breathing through the nose

Warnings
Do not use
■ to sedate a child or make a child sleepy
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's Disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this drug.

Ask a doctor before use if you have
■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are
■ taking any other nasal decongestant or stimulant ■ taking sedatives or tranquilizers

When using this product
Do not exceed recommended dosage.
■ marked drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children.

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur. ■ symptom do not improve within 7 days or are accompanied by fever.

200958-02 Rev. 06/2025

PEEL

LOHIST D

chlorpheniramine maleate / pseudoephedrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68047-120
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg in 5 mL
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
METHYLPARABEN (UNII: A2I8C7HI9T)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM ANHYDROUS (UNII: I4807BK602)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68047-120-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/15/2003	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/15/2003	

Labeler - Larken Laboratories, Inc. (149484540)

Registrant - Larken Laboratories, Inc. (149484540)

Revised: 10/2025

Larken Laboratories, Inc.