

CHLORMAX 50- chlortetracycline granule
Phibro Animal Health

ChlorMax®
50

ChlorMax 50

ChlorMax 50

Chlortetracycline
Type A Medicated Article
Antibacterial

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Active drug ingredient

Chlortetracycline (as hydrochloride) 50 g/lb

Inert ingredients

Roughage products, silicon dioxide, mineral oil

For use in manufacturing medicated feed only

Indications for Use

Indications for use	Chlortetracycline mg / lb body wt / day
Cattle Beef cattle (over 700 lbs): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline.	0.5
Withdraw 48 hours prior to slaughter. Calves, beef and non-lactating dairy cattle: Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to chlortetracycline. Treat for not more than 5 days.	10
Withdraw 24 hours prior to slaughter. Swine Control of porcine proliferative enteropathies (ileitis) caused by <i>Lawsonia intracellularis</i> susceptible to chlortetracycline. Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and	10

Salmonella choleraesuis and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to chlortetracycline. (Note: This drug level is equivalent to 400 grams per ton, depending on feed consumption and body weight).

Feed for not more than 14 days.

Turkeys

Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline. Feed continuously for 7 to 14 days.

25

Indications for use

Cattle

Growing cattle (over 400 lbs): For the reduction of the incidence of liver abscesses.

70

Beef cattle: For the control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella* spp. susceptible to chlortetracycline.

350

Withdraw 48 hours prior to slaughter.

Beef cattle (under 700 lbs): Control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline.

350

Withdraw 48 hours prior to slaughter.

Sheep

Breeding sheep: Reducing the incidence of (vibrionic) abortion caused by *Campylobacter fetus* infection susceptible to chlortetracycline.

80

Indications for use

Chickens

Control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to chlortetracycline. Feed continuously for 7 to 14 days.

100 - 200

Control of chronic respiratory disease (CRD) and air sac infection caused by *Mycoplasma gallisepticum* and *Escherichia coli* susceptible to chlortetracycline. Feed continuously for 7 to 14 days.

200 - 400

Reduction of mortality due to *Escherichia coli* infections susceptible to chlortetracycline.

500

Feed for 5 days.

Withdraw 24 hours prior to slaughter.

Turkeys

Control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to chlortetracycline.

200

Feed continuously for 7 to 14 days.

Control of hexamitiasis caused by *Hexamita meleagridis* susceptible to chlortetracycline. Feed continuously for 7 to

400

mg / head / day

In complete feed
Chlortetracycline g / ton

14 days.

Turkey pouls not over 4 weeks of age: Reduction of mortality due to paratyphoid caused by *Salmonella typhimurium* susceptible to chlortetracycline.

400

Swine

Reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by Group E Streptococci susceptible to chlortetracycline.

50 - 100

Breeding swine: Control of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by *Leptospira pomona* susceptible to chlortetracycline.

400

Feed continuously for not more than 14 days.

Mixing directions

For use in manufacture of complete feeds, mix thoroughly as follows:

Chlortetracycline, g/ton	ChlorMax 50, lb/ton
50 - 100	1 - 2
100 - 200	2 - 4
200 - 400	4 - 8
500	10

For use in manufacture of feed supplements to be fed at a rate of 1 lb per head per day, mix thoroughly as follows:

mg/lb body wt/day	If animal wt = 1000 lbs	ChlorMax 50 lb/ton supplement
0.5	500 mg	20
5.0	5,000 mg	200

Caution

For control and treatment of diseases caused by organisms susceptible to chlortetracycline as indicated. Observe the limitations of use and withdrawal periods as indicated.

Warning

Do not feed to chickens or turkeys producing eggs for human consumption.

A withdrawal period has not been established for this product in preruminating calves.

Do not use in calves to be processed for veal.

Storage

Store below 25°C (77°F), excursions permitted to 37°C (99°F).

Keep container tightly closed to prevent contamination or moisture pick-up.

For Use in Dry Feeds Only.

Not for Use in Liquid Medicated Feeds.

Restricted Drug (California) - Use only as directed.

Not for human use.

Approved by FDA under NADA # 046-699

Questions/Comments? For technical assistance or to obtain a Safety Data Sheet(s), contact Phibro Animal Health Corporation at 888-403-0074 or www.pahc.com. To report side effects, contact Phibro Animal Health Corporation at 1-888-475-7355. For additional information about reporting side effects, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

Phibro, Phibro logo and ChlorMax are trademarks owned by or licensed to Phibro Animal Health Corporation or its affiliates.

Distributed by:

Phibro Animal Health Corporation
Teaneck, NJ 07666

Net Wt 50 LB (22.68 kg)

176-9007-02


TAKE TIME



**OBSERVE LABEL
DIRECTIONS**

Label

ChlorMax 50



CHLORMAX 50

chlortetracycline granule

Product Information

Product Type	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:66104-1212
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Active Ingredient	Product Name	Description	Color
chlortetracycline	chlortetracycline	chlortetracycline	white
chlortetracycline	chlortetracycline	chlortetracycline	white
chlortetracycline	chlortetracycline	chlortetracycline	white

Ingredient Name	Basis of Strength	Strength
CHLORTETRACYCLINE (UNII: WCK1KIQ23Q) (CHLORTETRACYCLINE - UNII:WCK1KIQ23Q)	CHLORTETRACYCLINE	50 g in 0.45 kg

Product Characteristics

Color	brown (Light Tan to Brown)	Score	
Shape		Size	
Flavor			Imprint Code
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66104-1212-1	22.68 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA046699	02/15/2016	

Labeler - Phibro Animal Health (006989008)

Revised: 1/2026

Phibro Animal Health