

**GOOD SENSE MIGRAINE FORMULA- acetaminophen, aspirin, caffeine tablet, film coated**

**L. Perrigo Company**

-----  
**Perrigo Migraine Formula Drug Facts**

**Active ingredients (in each tablet)**

Acetaminophen 250 mg

Aspirin 250 mg (NSAID\*)

Caffeine 65 mg

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever

Pain reliever aid

**Use**

- treats migraine

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

you take

- more than 2 gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Medication overuse headache warning:** Headaches may worsen if this product is used for 10 or more days per month.

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause severe nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

### **Ask a doctor before use if**

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever

- reducers
- you have vomiting with migraine headache

### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for:
- diabetes
- gout
- arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not use more than directed**
- adults: take 2 gels with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 gels in 24 hours, unless directed by a doctor
- under 18 years of age: as a doctor

### **Other information**

- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

### **Inactive ingredients**

black iron oxide, carnauba wax, crospovidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, gelatin, glycerin, hypromellose, maltitol solution, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polyvinyl acetate phthalate, povidone, pregelatinized starch, propylene glycol, stearic acid, titanium dioxide

### **Questions or comments?**

**1-800-719-9260**

### **Package/Label Principal Display Panel**

GoodSense®

Treats Migraine

Migraine Formula

Acetaminophen, Aspirin (NSAID) and Caffeine Tablets

Pain Reliever/Pain Reliever Aid

80 Geltabs\*\*

\*\*Gelatin Coated Tablets

Compare to active ingredients of Excedrin® Migraine

GOODSENSE.

NDC 0113-1619-27

Treats Migraine

# Migraine Formula

Acetaminophen, Aspirin (NSAID) and Caffeine Tablets  
Pain Reliever/Pain Reliever Aid



80 Geltabs\*\* \*\*Gelatin Coated Tablets

Compare to active ingredients of  
Excedrin® Migraine



### Drug Facts

Active Ingredients (in each geltab)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)	Pain reliever
Caffeine 65 mg	Pain reliever aid

\*Nonsteroidal anti-inflammatory drug

**Use** ■ treats migraine

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior, weakness and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin redness ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:  
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
 ■ more than 2 geltabs in 24 hours, which is the maximum daily amount  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks a very day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  
 ■ are age 60 or older  
 ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ have 3 or more alcoholic drinks every day while using this product

■ Take more or for a longer time than directed

### Drug Facts (continued)

**Medication overuse headache warning:** Headaches may worsen if this product is used for 10 or more days per month. Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

**Do not use**  
 ■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

**Ask a doctor before use if**  
 ■ you have never had a migraine diagnosed by a health professional  
 ■ you have a headache that is different from your usual migraines  
 ■ you have the worst headache of your life  
 ■ you have fever and stiff neck  
 ■ you have headaches beginning after or caused by heat injury, exertion, coughing or heaving  
 ■ you experience your first headache after the age of 50  
 ■ you have daily headaches  
 ■ you have a migraine so severe as to require bedrest  
 ■ you have liver disease  
 ■ stomach bleeding warning applies to you  
 ■ you have a history of stomach problems, such as heartburn  
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
 ■ you are taking a diuretic  
 ■ you have asthma  
 ■ you have problems or serious side effects from taking pain relievers or fever reducers  
 ■ you have vomiting with your migraine headache

**Ask a doctor or pharmacist before use if you are**  
 ■ taking a prescription drug for:  
 ■ diabetes ■ gout ■ arthritis  
 ■ under a doctor's care for any serious condition  
 ■ taking any other drug  
 ■ taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

### Drug Facts (continued)

**Stop use and ask a doctor if**  
 ■ an allergic reaction occurs. Seek medical help right away.  
 ■ you experience any of the following signs of stomach bleeding:  
 ■ feel faint ■ vomit blood ■ have bloody or black stools  
 ■ have stomach pain that does not get better  
 ■ your migraine is not relieved or worsens after the first dose  
 ■ new or unusual symptoms occur  
 ■ ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
 ■ do not use more than directed  
 ■ adults: take 2 geltabs with a glass of water  
 ■ if symptoms persist or worsen, ask your doctor  
 ■ do not take more than 2 geltabs in 24 hours, unless directed by a doctor  
 ■ under 18 years of age: ask a doctor

**Other information**  
 ■ store at 20°-25°C (68°-77°F) ■ close cap tightly after use  
 ■ read all product information before using. Keep this box for important information.

**Inactive ingredients:** black iron oxide, croscellose, croscellose, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, gelatin, glycerin, hydroxypropyl methylcellulose, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polyvinyl acetate, polyethylene glycol, polyethylene glycol, stearic acid, titanium dioxide

**Questions or comments?** 1-800-719-9260

KEEP BOX FOR REFERENCE  
 DO NOT USE IF RAINED FOR UNBLENDED  
 IS BROKEN OR IN SHAKEN

80 Geltabs

**Migraine Formula**

distributed by  
**Perrigo®**  
 Allegheny, PA 15001

100% Recycled Paper Box  
 100% Recycled Plastic Bottle

**GOOD SENSE MIGRAINE FORMULA**  
acetaminophen, aspirin, caffeine tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0113-1619
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>FERROSO FERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>CROSPROVIDONE</b> (UNII: 2S7830E561)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MALTITOL</b> (UNII: D65DG142WK)	
<b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYVINYL ACETATE PHTHALATE</b> (UNII: 58QVG85GW3)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	GREEN, WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	13mm
<b>Flavor</b>		<b>Imprint Code</b>	M
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0113-1619-27	1 in 1 CARTON	06/24/2022	
1		80 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075794	06/24/2022	

**Labeler** - L. Perrigo Company (006013346)

Revised: 6/2022

L. Perrigo Company