

**VITACILINA FIRST AID ANTIBIOTIC- bacitracin zinc, neomycin sulfate,  
polymyxin b sulfate ointment  
Taisho Pharmaceutical California Inc.**

-----  
**Vitacilina First Aid Antibiotic**

***Drug Facts***

***Active ingredient (in each gram)***

Bacitracin Zinc 400 units

Neomycin sulfate 3.5 mg

Polymyxin B sulfate 5,000 units

***Purpose***

First aid antibiotic

***Use***

First aid to help prevent infection in minor cuts, scrapes, and burns

***Warnings***

**For external use only.**

**Do not use**

- if you are allergic to any of the ingredients
- in the eyes
- over large areas of the body

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

**Stop use and ask a doctor if**

- you need to use longer than 1 week
- condition persists or gets worse
- rash or other allergic reaction develops

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center

**Directions**

- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage


**Other information**

- store at 15 to 30°C (59 to 86°F )
- DO NOT USE IF SEAL ON TUBE IS BROKEN OR MISSING. **Tamper Evident:**

**Inactive ingredient**

petrolatum

**Package Labeling 28g**



Add image transcription here...

FIRST AID ANTIBIOTIC

NDC 81929-003-01

# VITACILINA®

NET WT 1 OZ (28 g)

<b>Drug Facts</b>		<b>Drug Facts</b> (continued)	
<b>Active ingredient (in each gram)</b>	<b>Purpose</b>	<b>Stop use and ask a doctor if</b> ■ you need to use longer than 1 week ■ condition persists or gets worse ■ rash or other allergic reaction develops	
Bacitracin zinc 400 units.....	First aid antibiotic	<b>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center</b>	
Neomycin sulfate 3.5 mg.....	First aid antibiotic	<b>Directions</b> ■ clean the affected area ■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily ■ may be covered with a sterile bandage	
Polymyxin B sulfate 5,000 units....	First aid antibiotic	<b>Other information</b> ■ store at 15 to 30°C (59 to 86°F) ■ <b>Tamper Evident: DO NOT USE IF SEAL ON TUBE IS BROKEN OR MISSING.</b>	
<b>Use</b>	First aid to help prevent infection in minor cuts, scrapes, and burns	<b>Inactive ingredient</b> petrolatum	
<b>Warnings</b>	For external use only.	<b>DISTRIBUTED BY</b> Taisho Pharmaceutical California Inc. PL33103T 3528 Torrance Blvd, Suite #320 Torrance, CA 90503 U.S.A. V01	
<b>Do not use</b>	■ if you are allergic to any of the ingredients ■ in the eyes ■ over large areas of the body		
<b>Ask a doctor before use if you have</b>	■ deep or puncture wounds ■ animal bites ■ serious burns ▶		

Add image transcription here...

## Package Labeling 14g

NO  
COLOR

**Drug Facts** (continued)

**Other information** ■ store at 15 to 30°C (59 to 86°F)  
■ **Tamper Evident: DO NOT USE IF FLAPS ON CARTON ARE NOT SEALED OR MISSING**

**Inactive ingredient** petrolatum



PL33104B  
V01  
NO  
COLOR

**DISTRIBUTED BY**  
Taisho Pharmaceutical California Inc. 3528 Torrance Blvd, Suite #320 Torrance, CA 90503 U.S.A.



VITACILINA

FIRST AID ANTIBIOTIC

NDC 81929-003-02

**VITACILINA**<sup>®</sup>

NET WT 0.5 OZ (14 g)

NO  
COLOR

FIRST AID ANTIBIOTIC

NDC 81929-003-02

**VITACILINA**<sup>®</sup>

NET WT 0.5 OZ (14 g)

NO  
COLOR

GLUE FLAP  
NO COLOR / TEXT  
NO COATING

**Drug Facts**

<b>Active ingredient (in each gram)</b>	<b>Purpose</b>
Bacitracin zinc 400 units.....	First aid antibiotic
Neomycin sulfate 3.5 mg.....	First aid antibiotic
Polymyxin B sulfate 5,000 units.....	First aid antibiotic

**Use** First aid to help prevent infection in minor cuts, scrapes, and burns

**Warnings**

For external use only.

**Do not use** ■ if you are allergic to any of the ingredients  
■ in the eyes ■ over large areas of the body

**Drug Facts** (continued)

**Ask a doctor before use if you have** ■ deep or puncture wounds ■ animal bites ■ serious burns

**Stop use and ask a doctor if** ■ you need to use longer than 1 week ■ condition persists or gets worse ■ rash or other allergic reaction develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away: 800-222-1222.

**Directions** ■ clean the affected area ■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily  
■ may be covered with a sterile bandage

GLUE FLAP  
NO COLOR / TEXT  
NO COATING

FIRST AID ANTIBIOTIC

**VITACILINA®**

NET WT 0.5 OZ (14 g)

<b>Drug Facts</b>		<b>Drug Facts</b> (continued)
<b>Active ingredient (in each gram)</b>	<b>Purpose</b>	gets worse ■ rash or other allergic reaction develops
Bacitracin zinc 400 units.....	First aid antibiotic	<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away: 800-222-1222.
Neomycin sulfate 3.5 mg.....	First aid antibiotic	
Polymyxin B sulfate 5,000 units..	First aid antibiotic	
<b>Use</b>	First aid to help prevent infection in minor cuts, scrapes, and burns	<b>Directions</b> ■ clean the affected area ■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area up to 3 times daily ■ may be covered with a sterile bandage
<b>Warnings</b> For external use only.		<b>Other information</b> ■ store at 15 to 30°C (59 to 86°F)
<b>Do not use</b> ■ if you are allergic to any of the ingredients ■ in the eyes ■ over large areas of the body		<b>Inactive ingredient</b> petrolatum
<b>Ask a doctor before use if you have</b> ■ deep or puncture wounds ■ animal bites ■ serious burns		<b>DISTRIBUTED BY</b> NDC 81929-003-02 Taisho Pharmaceutical California Inc. 3528 Torrance Blvd, Suite #320 Torrance, CA 90503 U.S.A. PL33104T V01
<b>Stop use and ask a doctor if</b> ■ you need to use longer than 1 week ■ condition persists or ►		

## VITACILINA FIRST AID ANTIBIOTIC

bacitracin zinc, neomycin sulfate, polymyxin b sulfate ointment

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:81929-003
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BACITRACIN ZINC</b> (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	400 U in 1 g
<b>NEOMYCIN SULFATE</b> (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 g
<b>POLYMYXIN B SULFATE</b> (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	5000 U in 1 g

### Inactive Ingredients

Ingredient Name		Strength		
PETROLATUM (UNII: 4T6H12BN9U)				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81929-003-01	1 in 1 CARTON	07/01/2022	01/26/2028
1		28 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:81929-003-02	1 in 1 CARTON	07/01/2022	04/29/2027
2		14 g in 1 TUBE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M004	07/01/2022	01/26/2028	

**Labeler** - Taisho Pharmaceutical California Inc. (603827635)

Revised: 8/2025

Taisho Pharmaceutical California Inc.