

WOMENS GENTLE LAXATIVE- bisacodyl tablet, delayed release
Chain Drug Consortium

Premier Value 44-676A

Active ingredient (in each tablet)

Bisacodyl USP, 5 mg

Purpose

Stimulant laxative

Uses

- for relief of occasional constipation (irregularity)
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- you may have stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take with a glass of water

adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years	take 1 tablet in a single daily dose
children under 6 years	ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- avoid excessive humidity
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

carmine, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal display panel

**Premier
Value®**

**Women's
Gentle Laxative**

Bisacodyl USP, 5 mg

STIMULANT LAXATIVE

Gentle, dependable
constipation relief

30 Tablets

Enteric coated tablets

actual size

INDEPENDENTLY TESTED

PV

SATISFACTION GUARANTEED

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

50844 REV0119A67601

**Distributed By:
Pharmacy Value Alliance, LLC
407 East Lancaster Avenue,
Wayne, PA 19087**

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.



TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Drug Facts

Active ingredient (in each tablet)

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Purpose

Bisacodyl USP, 5 mg Stimulant laxative

Uses

■ for relief of occasional constipation (irregularly) ■ this product generally produces bowel movement in 6 to 12 hours

Drug Facts (continued)



Women's Gentle Laxative

Bisacodyl USP, 5 mg

STIMULANT LAXATIVE

Gentle, dependable constipation relief



actual size

30 Tablets

Enteric coated tablets



no print/ no varnish area lot/no. & exp. date



8 40986 03226 4

Drug Facts (continued)

polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-426-9391

Distributed By:
Pharmacy Value Alliance, LLC
407 East Lancaster Avenue,
Wayne, PA 19087

B-1590-676A-01
REV0119A67601

50844 REV0119A67601

Drug Facts (continued)

Warnings
Do not use if you cannot swallow without chewing.
Ask a doctor before use if you have a sudden change in bowel habits that lasts more than 2 weeks ■ stomach pain, nausea or vomiting

When using this product
■ do not chew or crush tablet(s)
■ do not use within 1 hour after taking an antacid or milk

■ you may have stomach discomfort, flatness and cramps

Stop use and ask a doctor if
■ you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.

■ you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions
■ take with a glass of water
adults and children 12 years and over take 1 to 3 tablets in a single daily dose
children 6 to under 12 years take 1 tablet in a single daily dose
children under 6 years ask a doctor

Other information

TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
■ avoid excessive humidity
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ see end flap for expiration date and lot number

Inactive ingredients camphine, colloidal silicon dioxide, corn starch, FD & C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcrystalline cellulose, polydextrose,

Premier Value 44-676A

WOMENS GENTLE LAXATIVE

bisacodyl tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68016-197
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B71O)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	B
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-197-30	2 in 1 CARTON	09/16/2015	
1		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	09/16/2015	

Labeler - Chain Drug Consortium (101668460)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(68016-197)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(68016-197) , pack(68016-197)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(68016-197)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(68016-197)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(68016-197)

Revised: 9/2023

Chain Drug Consortium