

ASPIRIN 81MG- aspirin tablet
Innovida Pharmaceutique Corporation

Active Ingredient

Aspirin 81 mg(NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains
- ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you are taking a diuretic
- you have a history of stomach problems, such as heartburn
- you have: -high blood pressure -heart disease -liver cirrhosis -kidney disease -asthma

Ask a doctor or pharmacist before use if you are

- taking any other drug containing an NSAID (prescription or nonprescription)

- taking a blood thinning (anticoagulant) or steroid drug
- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: -feel faint -vomit blood
- have bloody or black stools -have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- any new symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. it is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- swallow whole, do not chew or crush
- do not exceed recommended dose
- adults and children 12 years and older: take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at controlled room temperature

Inactive Ingredients

cellulose, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, PEG, polydextrose, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, starch, talc, titanium dioxide, triacetin, triethyl citrate, wax.

Questions or Comments? 1-888-462-4166

Product label

NDC: 71800-040-02

INNNOVIDA
PHARMACEUTICAL CORPORATION

Aspirin

Pain Reliever (NSAID),
81 mg

Enteric Coated Tablet

250 Tablets

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet)	Purpose
Aspirin 81 mg(NSAID)*	Pain reliever
<small>*nonsteroidal anti-inflammatory drug</small>	

Uses • for the temporary relief of minor aches and pains • ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if • the stomach bleeding warning applies to you • you are taking a diuretic • you have a history of stomach problems, such as heartburn • you have: • high blood pressure • heart disease • liver cirrhosis • kidney disease • asthma

Mfd. for:
Innovida Pharmaceutique
Corporation, Phoenix, AZ 85040
Tel.: 888-462-4166



PEEL HERE FOR MORE DRUG FACTS

Questions or Comments? 1-888-462-4166

Inactive ingredients: cellulose, D&C yellow #10, FD & C yellow #6, hypromellose, methylacrylic acid, Fe₃O₄, polydextrose, silica, stearic acid, sodium bicarbonate, sodium lauryl sulfate, starch, talc, titanium dioxide, triacetin, triethyl citrate, wax

Other information • store at controlled room temperature

12 years; ask a doctor

tablets in 24 hours, or as directed by a doctor • children under 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours • adults and children 12 years and older: swallow whole, do not chew or crush • do not take with other aspirin products • drink a full glass of water with each dose

In case of overdose, get medical help or contact Poison Control Center right away.

Keep out of reach of children.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless directly directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Stop use and ask a doctor if • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have blood or black stools • have stomach pain that does not get better • an allergic reaction occurs. Seek medical help if • pain gets worse or lasts more than 10 days • redness or swelling is present • fever, or hives or rash • more than 3 days any new symptoms occur • ringing in the ears or loss of hearing occurs

Ask a doctor or pharmacist before use if you are • taking any other drug containing an NSAID (prescription or nonprescription) • taking a blood thinning (anticoagulant) or steroid drug • taking aspirin drug or diuretic, gout or arthritis

Drug Facts (continued)

ASPIRIN 81MG

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71800-040
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ACETYLSALICYLIC ACID - UNII:R16CO5Y76E)		ASPIRIN	81 mg	
Inactive Ingredients				
Ingredient Name			Strength	
STARCH, CORN (UNII: O8232NY3SJ)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
HYPROMELLOSES (UNII: 3NXW29V3WO)				
METHACRYLIC ACID (UNII: 1CS02G8656)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
POLYDEXTROSE (UNII: VH2XOU12IE)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
SHELLAC (UNII: 46N107B710)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
DIMETHICONE (UNII: 92RU3N3Y10)				
SODIUM BICARBONATE (UNII: 8MDF5V39QO)				
SODIUM LAURYL SULFATE (UNII: 368GB5141J)				
TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
TRIACETIN (UNII: XHX3C3X673)				
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)				
Product Characteristics				
Color	yellow	Score	no score	
Shape	ROUND	Size	6mm	
Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71800-040-02	250 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/2025	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	02/10/2025		

Labeler - Innovida Pharmaceutique Corporation (080892908)