

CAREONE ALLERGY RELIEF D- cetirizine hcl, pseudoephedrine hcl tablet, extended release
American Sales Company

American Sales Company Allergy Relief-D Drug Facts

Active ingredients (in each extended release tablet)

Cetirizine HCl 5 mg

Pseudoephedrine HCl 120 mg

Purpose

Antihistamine

Nasal Decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- diabetes
- glaucoma
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are

taking tranquilizers or sedatives.

When using this product

- do not use more than directed
- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- you get nervous, dizzy, or sleepless
- symptoms do not improve within 7 days or are accompanied by fever

If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not break or chew tablet; swallow tablet whole

adults and children 12 years and over	take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
adults 65 years and over	ask a doctor
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- store between 20° to 25°C (68° to 77°F)

Inactive ingredients

colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, talc, yellow iron oxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredients in Zyrtec-D®

ORIGINAL PRESCRIPTION STRENGTH

12hr

Allergy Relief-D

Cetirizine Hydrochloride and Pseudoephedrine Hydrochloride Extended Release Tablets, 5 mg/120 mg

Antihistamine/Nasal Decongestant

Actual Size

24 EXTENDED RELEASE TABLETS

Allergy & Congestion

INDOOR & OUTDOOR ALLERGIES

OUR PHARMACISTS RECOMMEND

12-HOUR RELIEF OF:

Sneezing – Runny Nose – Sinus Pressure

Itchy, Watery Eyes – Itchy Throat or Nose – Nasal Congestion

Gluten Free



Drug Facts (continued)
Questions or comments? 1-800-719-9260
 Made in India
 DISTRIBUTED BY FOCUS PHARMACEUTICALS, LLC
 LANDOVER, MD 20785 • 1-877-846-9949
 ©2017 S&S Brands, LLC
 Quality guaranteed or your money back.

Drug Facts (continued)

Warnings
 ■ If you have ever had an allergic reaction to this product or any of its ingredients or to an anti-histamine containing hydroxyzine.
 ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or remittal conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if you are taking a MAOI drug, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
 ■ heart disease ■ thyroid disease ■ diabetes ■ glaucoma ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ liver or kidney disease. Your doctor should determine if you need a different dose.
Ask a doctor or pharmacist before use if you are taking
 tranquilizers or sedatives.
When using this product
 ■ do not use more than directed ■ avoid alcoholic drinks ■ do not use with other MAOIs ■ avoid alcohol, sedatives, and tranquilizers as they may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery
 ■ an allergic reaction to this product occurs.
 ■ Seek medical help if right away.

Directions
 ■ Do not break or chew tablet; swallow tablet whole
 ■ Adults and children: take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
 ■ Adults 65 years and over: ask a doctor
 ■ Children under 12 years of age: ask a doctor
 ■ Ask a doctor or pharmacist for information on liver or kidney disease.

Other information
 ■ Store between 20° to 25°C (68° to 77°F)

Inactive ingredients
 colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, talc, yellow iron oxide

Drug Facts (continued)

Warnings
 ■ You get nervous, dizzy, or sleepless
 ■ Symptoms do not improve within 7 days or are accompanied by fever
 ■ If pregnant or breastfeeding: not recommended
 ■ If breastfeeding: not recommended
 ■ If pregnant: ask a health professional before use.
 ■ Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ Do not break or chew tablet; swallow tablet whole
 ■ Adults and children: take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
 ■ Adults 65 years and over: ask a doctor
 ■ Children under 12 years of age: ask a doctor
 ■ Ask a doctor or pharmacist for information on liver or kidney disease.

Other information
 ■ Store between 20° to 25°C (68° to 77°F)

Inactive ingredients
 colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, talc, yellow iron oxide

DO NOT USE IF BUSTER UNIT IS BROKEN OR TORN
 *This product is not manufactured or distributed by McKellicon Consumer Healthcare, division of McNeil-PPC, Inc., distributor of Zyrtec-D®.

Drug Facts (continued)

Warnings
 ■ If pregnant or breastfeeding: not recommended
 ■ If breastfeeding: not recommended
 ■ If pregnant: ask a health professional before use.
 ■ Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ Do not break or chew tablet; swallow tablet whole
 ■ Adults and children: take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
 ■ Adults 65 years and over: ask a doctor
 ■ Children under 12 years of age: ask a doctor
 ■ Ask a doctor or pharmacist for information on liver or kidney disease.

Other information
 ■ Store between 20° to 25°C (68° to 77°F)

Inactive ingredients
 colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, talc, yellow iron oxide

Drug Facts (continued)

Warnings
 ■ If pregnant or breastfeeding: not recommended
 ■ If breastfeeding: not recommended
 ■ If pregnant: ask a health professional before use.
 ■ Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ Do not break or chew tablet; swallow tablet whole
 ■ Adults and children: take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
 ■ Adults 65 years and over: ask a doctor
 ■ Children under 12 years of age: ask a doctor
 ■ Ask a doctor or pharmacist for information on liver or kidney disease.

Other information
 ■ Store between 20° to 25°C (68° to 77°F)

Inactive ingredients
 colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, talc, yellow iron oxide

CAREone®
 Compare to the active ingredients in Zyrtec-D®*
 ORIGINAL PRESCRIPTION STRENGTH
12hr
Allergy Relief-D
 Cetirizine Hydrochloride and Pseudoephedrine Hydrochloride Extended Release Tablets, 5 mg/120 mg Antihistamine/Nasal Decongestant



Actual Size

24
EXTENDED RELEASE TABLETS
 Allergy & Congestion
 INDOOR & OUTDOOR ALLERGIES

17462 OF C1



12-HOUR RELIEF OF:

• Sneezing • Runny Nose • Sinus Pressure
• Itchy, Watery Eyes • Itchy Throat or Nose • Nasal Congestion

Gluten Free

CAREONE ALLERGY RELIEF D

cetirizine hcl, pseudoephedrine hcl tablet, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-091
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CETIRIZINE HYDROCHLORIDE (UNII: 640047KTOA) (CETIRIZINE - UNII:YO7261ME24)	CETIRIZINE HYDROCHLORIDE	5 mg
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (160000 WAMW) (UNII: RFW2ET671P)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
TALC (UNII: 7SEV7J4R1U)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	WHITE (one side white one side light yellow)	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	5029;5;120
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-091-62	24 in 1 CARTON	07/20/2016	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077170	07/20/2016	

Labeler - American Sales Company (809183973)

Revised: 12/2019

American Sales Company