

**ALEVE CAPLETS - EASY OPEN ARTHRITIS CAP- naproxen sodium tablet
Bayer HealthCare LLC.**

Aleve ®

Caplets - Soft Grip Arthritis Cap

Drug Facts

Active ingredient (in each caplet)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID) ¹

¹ nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Allergy alert

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance

is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body

- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

-
- | | |
|--|--|
| <ul style="list-style-type: none"> • Adults and children 12 years and older | <ul style="list-style-type: none"> • take 1 tablet every 8 to 12 hours while symptoms last • for the first dose you may take 2 tablets within the first hour • do not exceed 2 tablets in any 8- to 12-hour period • do not exceed 3 tablets in a 24-hour period |
| <ul style="list-style-type: none"> • Children under 12 years | <ul style="list-style-type: none"> • ask a doctor |
-

Other information

- **each caplet contains:** sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients

FD&C blue #2 lake, hypromelloses, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidones, talc, titanium dioxide

Questions or comments?

1-800-395-0689 (Mon - Fri 9AM - 5PM EST)

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981

PRINCIPAL DISPLAY PANEL - 270 Caplet Bottle Carton

SOFT GRIP ARTHRITIS CAP

ALL DAY STRONG®

ALEVE®

naproxen sodium tablets, 220 mg (NSAID)

Pain reliever/fever reducer

THIS PACKAGE IS CHILD RESISTANT

STRENGTH TO LAST 12 HOURS

ACTUAL SIZE

270 CAPLETS

CAPSULE-SHAPED TABLETS



ALEVE CAPLETS - EASY OPEN ARTHRITIS CAP

naproxen sodium tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0280-6050
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL (capsule shaped tablets)	Size	12mm
Flavor		Imprint Code	Aleve
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-6050-50	1 in 1 CARTON	11/11/2013	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0280-6050-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/11/2013	
3	NDC:0280-6050-27	1 in 1 CARTON	11/11/2013	
3		270 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0280-6050-02	24 in 1 BOTTLE; Type 0: Not a Combination Product	06/11/2020	
5	NDC:0280-6050-03	1 in 1 CARTON	11/11/2013	
5		130 in 1 BOTTLE; Type 0: Not a Combination		

5		Product	
6	NDC:0280-6050-04	1 in 1 CARTON	11/11/2013
6		150 in 1 BOTTLE; Type 0: Not a Combination Product	
7	NDC:0280-6050-06	90 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2021



Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020204	11/11/2013	

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 10/2022

Bayer HealthCare LLC.