# ALEVE CAPLETS - EASY OPEN ARTHRITIS CAP- naproxen sodium tablet Bayer HealthCare LLC.

-----

Aleve ®

Caplets - Soft Grip Arthritis Cap

**Drug Facts** 

## Active ingredient (in each caplet)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID) <sup>1</sup>

1 nonsteroidal anti-inflammatory drug

## **Purposes**

Pain reliever/fever reducer

#### Uses

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - menstrual cramps
  - headache
  - toothache
  - the common cold
- temporarily reduces fever

## Allergy alert

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

## Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance

is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

## Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

## Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

## When using this product

take with food or milk if stomach upset occurs

## Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body

- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

## If pregnant or breast-feeding

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

#### Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul> <li>take 1 tablet every 8 to 12 hours while symptoms last</li> <li>for the first dose you may take 2 tablets within the first hour</li> <li>do not exceed 2 tablets in any 8- to 12-hour period</li> <li>do not exceed 3 tablets in a 24-hour period</li> </ul>
Children under 12 years	ask a doctor

#### Other information

- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F)

## **Inactive ingredients**

FD&C blue #2 lake, hypromelloses, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidones, talc, titanium dioxide

#### Questions or comments?

**1-800-395-0689** (Mon - Fri 9AM - 5PM EST)

Dist. by: Bayer HealthCare LLC Whippany, NJ 07981

## PRINCIPAL DISPLAY PANEL - 270 Caplet Bottle Carton

SOFT GRIP ARTHRITIS CAP

ALL DAY STRONG ®
ALEVE ®
naproxen sodium tablets, 220 mg (NSAID)
Pain reliever/fever reducer


THIS PACKAGE IS CHILD RESISTANT

STRENGTH TO LAST 12 HOURS

**ACTUAL SIZE** 

270 CAPLETS

**CAPSULE-SHAPED TABLETS** 



#### **ALEVE CAPLETS - EASY OPEN ARTHRITIS CAP**

naproxen sodium tablet

<b>Product</b>	Inform	ation
PIUUULL		aliui

**Product Type** HUMAN OTC DRUG **Item Code (Source)** NDC:0280-6050

**Route of Administration** ORAL

## **Active Ingredient/Active Moiety**

1	,		
	Ingredient Name	<b>Basis of Strength</b>	Strength
	NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

## **Inactive Ingredients Ingredient Name**

FD&C BLUE NO. 2 (UNII: L06K8R7DQK) HYPROMELLOSES (UNII: 3NXW29V3WO) MAGNESIUM STEARATE (UNII: 70097M6I30) CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3MJQ0SDW1A)

Strength

POVIDONE (UNII: FZ989GH94E)

TALC (UNII: 7SEV7J4R1U)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

## **Product Characteristics**

Color	blue	Score	no score
Shape	OVAL (capsule shaped tablets)	Size	12mm
Flavor		Imprint Code	Aleve
Contains			

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:0280-6050- 50	1 in 1 CARTON	11/11/2013			
1		50 in 1 BOTTLE; Type 0: Not a Combination Product				
2	NDC:0280-6050- 01	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/11/2013			
3	NDC:0280-6050- 27	1 in 1 CARTON	11/11/2013			
3		270 in 1 BOTTLE; Type 0: Not a Combination Product				
4	NDC:0280-6050- 02	24 in 1 BOTTLE; Type 0: Not a Combination Product	06/11/2020			
5	NDC:0280-6050- 03	1 in 1 CARTON	11/11/2013			
_		130 in 1 BOTTLE; Type 0: Not a Combination				

3		Product					
6	NDC:0280-6050- 04	1 in 1 CARTON		-	11/11/2013		
6		150 in 1 BOTTLE; Ty Product	ype 0: Not a Combin	nation			
7	NDC:0280-6050- 06	90 in 1 BOTTLE; Typ Product	oe 0: Not a Combina	ation	07/01/2021		
8	NDC:0280-6050- 05	110 in 1 BOTTLE; Ty Product	/pe 0: Not a Combin	nation	07/01/2021		
	de mariant de constant de la constan	ingen of security blacking them the control graphers (see the control graphers (see the control graphers (see the control graphers (see the control graphers) of security of s	NGTH AST 270 CAPIETS OF STATES OF ST	A approximately soft on process so dis Point solitones/for	ARTHRITIS CAP  ALL DAY STROMO.  LEVE  Middle, 220 mg (NSAID)  WALK IN DECEMBER	Dong Facility Antique (Sept of Sept of	construction analysis alone to the desired to the construction of the landers of the construction of the landers of the lander
M		nformation					
	Marketing Category		Number or Mond Citation	ograph	Marketing S Date	Start	Marketing End Date
NΕ	DA	NDA020204			11/11/2013		

## **Labeler -** Bayer HealthCare LLC. (112117283)

Revised: 4/2025 Bayer HealthCare LLC.