# PRIMATENE MIST- epinephrine inhalation aerosol Armstrong Pharmaceuticals, Inc.

-----

# Active ingredient (in each spray)

Epinephrine 0.125 mg

## **Purpose**

Bronchodilator

#### Uses

for temporary relief of mild symptoms of intermittent asthma

- wheezing
- tightness of chest
- shortness of breath

## Warnings

# For oral inhalation only

Asthma alert: Because asthma may be life threatening, see a doctor if you

- are not better in 20 minutes
- get worse
- need more than 8 inhalations in 24 hours
- have more than 2 asthma attacks in a week

These may be signs that your asthma is getting worse.

#### Do not use

- unless a doctor said you have asthma
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs taken for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or a pharmacist before taking this product.

# Ask a doctor before use if you have

- ever been hospitalized for asthma
- heart disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- thyroid disease
- seizures
- narrow angle glaucoma

# Ask a doctor or pharmacist before use if you are

- taking prescription drugs for asthma, obesity, weight control, depression, or psychiatric or emotional conditions
- taking any drug that contains phenylephrine, pseudoephedrine, ephedrine, or caffeine (such as for allergy, cough-cold, or pain)

# When using this product

- your blood pressure or heart rate may go up. This could increase your risk of heart attack or stroke, which may cause death.
- **■** your risk of heart attack or stroke increases if you:
  - have a history of high blood pressure or heart disease
- take this product more frequently or take more than the recommended dose.
- avoid foods or beverages that contain caffeine
- avoid dietary supplements containing ingredients reported or claimed to have a stimulant effect
- avoid spraying in eyes
- contents under pressure. Do not puncture or incinerate.
- do not store near open flame or heat above 120°F (49°C). May cause bursting.

# Stop use and ask a doctor if

- your asthma is getting worse (see asthma alert)
- you have difficulty sleeping
- you have a rapid heart beat
- you have tremors, nervousness, or seizure

If pregnant or breast-feeding, ask a health care professional before use

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

■ read the Consumer Information Insert for detailed directions on how to use this product

#### ■ do not use more than directed

- for adults and children 12 years of age and over
- children under 12 years of age: do not use; it is not known if the drug works or is safe in children under 12

# Before first use (new inhaler)

activate new inhaler by shaking then spraying into air 4 separate times

# each time you dose

- remove red cap (if attached)
- shake then spray into the air 1 time
- exhale completely, place inhaler in mouth
- inhale deeply while pressing down on top of inhaler, then continue the deep breath
- hold breath as long as possible, exhale

wait 1 minute. If symptoms not relieved, take a second inhalation by repeating steps above.

#### after use

- wait at least 4 hours between doses
- do not use more than 8 inhalations in 24 hours
- wash inhaler after each day of use. Run water through the mouthpiece for 30 seconds.

#### Other information

- store at room temperature, between 15-25°C (59-77°F)
- contains no sulfites
- keep this label and enclosed materials. They contain important additional information.

### Inactive ingredients

dehydrated alcohol (1%), hydrofluoroalkane (HFA 134a), polysorbate 80, thymol

#### Questions or comments?

call 1-8 Primatene or 1-877-462-8363

Mon.- Fri. 7 am - 5 pm PST

**Primatene MIST® Carton** 

**Principal Display Panel Text:** 

**Primatene MIST®** 

Epinephrine Inhalation Aerosol 0.125 mg per spray Bronchodilator For Oral Inhalation Only

For TEMPORARY relief of MILD symptoms of INTERMITTENT asthma

Suspension:

**Shake then Spray** into the air before each inhalation.

This item is


electronically

protected

**NEW FORMULATION:** 

# See Important Usage Information on Insert and on Side Panels

160 Metered Sprays Net Weight: 11.7g



# **PRIMATENE MIST**

epinephrine inhalation aerosol

raa	li ict	Intorn	nation
IUU	IULL		nation

Product Type HUMAN OTC DRUG Item Code (Source) NDC:17270-5530

Route of Administration RESPIRATORY (INHALATION)

# **Active Ingredient/Active Moiety**

Ingredient Name	<b>Basis of Strength</b>	Strength
EPINEPHRINE (LINII: YKH83404RH) (EPINEPHRINE - LINII: YKH83404RH)	FPINEPHRINE	125 ug

Inactive	Ingredients

Ingredient Name	Strength

ALCOHOL (UNII: 3K9958V90M)

NORFLURANE (UNII: DH9E53K1Y8)

POLYSORBATE 80 (UNII: 60ZP39ZG8H)	
THYMOL (UNII: 3I50XA376E)	

Packaging				
#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17270- 5530-0	1 in 1 CARTON	12/14/2018	
1		160 in 1 INHALER; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA	NDA205920	12/14/2018		

# Labeler - Armstrong Pharmaceuticals, Inc. (809773794)

Establishment				
Name	Address	ID/FEI	Business Operations	
Armstrong Pharmaceuticals, Inc.		809773794	manufacture(17270-5530), analysis(17270-5530)	

Revised: 7/2020 Armstrong Pharmaceuticals, Inc.