# COUGH DM- dextromethorphan polistirex suspension Kroger Company

-----

## Kroger Co. Cough DM Drug Facts

## Active ingredient (in each 5 mL)

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide

## Purpose

Cough suppressant

### Uses

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

## Warnings

### Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Allergy Alert:** Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

## Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

## Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## **Directions**

- shake bottle well before use
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by doctor

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24
	hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24
	hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24
	hours
children under 4 years of age	do not use

## Other information

- each 5 mL contains: sodium 5 mg
- store at  $20^{\circ}$  to  $25^{\circ}$ C ( $68^{\circ}$  to  $77^{\circ}$ F)
- dosing cup provided

## **Inactive Ingredients**

D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, glycerin, high fructose corn syrup, methylparaben, natural and artificial orange flavor, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

### Questions or comments?

1-800-632-6900

## Package/Label Principal Display Panel

COMPARE TO the active ingredient of DELSYM® ORANGE-FLAVOR

See side panel

OUR PHARMACIST RECOMMENDED

Cough DM

Dextromethorphan Polistirex Extended-Release Oral Suspension

COUGH SUPPRESSANT

Contains Sodium Metabisulfite, a Sulfite That May Cause Allergic-Type Reactions

Day or Night

Alcohol-Free

Dosing Cup Included

12 HOUR COUGH RELIEF

Orange-Flavored Liquid

3 FL OZ (89 mL)



#### Drug Facts

Active ingredient (in each 5 mL) Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide....

Purpose

Uses temporarily relieves
■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants

■ the impulse to cough to help you get to sleep

#### Warnings

Warmings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, Ifyou do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert Contains sodium metabisulfite, a suffite that may cause

allergic-type reactions.

allergic-type reactions.

Ask a doctor before use if you have

chronic cough that lasts as occurs with smoking, asthma or emphysema

cough that occurs with too much phlegm (mucus)

■ cough that occurs with too much philegm (mucus)
Stop use and sak a doctor if
■ side effects occur. You may report side effects to FIDA at 1-800-FIDA-1088.
■ cough lasts more than 7 days, cough comes back, or occurs with fever, rash
or headach e that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdoes, get medical help or
contact a Poison Control Center right away (1-800-222-1222).

#### Directions

#### ■ shake bottle well before use

- measure only with dosing cup provided
   do not use dosing cup with other products
   dose as follows or as directed by a doctor

adults and children	10 mL every 12 hours,
12 years of age and over	not to exceed 20 mL in 24 hours
children 6 to under	5 mL every 12 hours,
12 years of age	not to exceed 10 mL in 24 hours
children 4 to under	2.5 mL every 12 hours,
6 years of age	not to exceed 5 mL in 24 hours
171 1 1 1	1 .

#### Other information

■ each 5 mL contains: sodium 5 mg ■ store at 20° to 25°C (68° to 77°F) ■ dosing cup provided

#### Drug Facts (continued)

Drug Facts (CORTINUEG)

Inactive ingredients D&C Red #30
aluminum lake, D&C Yellow #10 aluminum lake,
glycerin, high fructose com syrup, methylparaben,
natural and artificial orange flavor, polysorbate 80,
polyvinyl acetate, povidone, propylparaben,
purified water, sodium metabisulifite, sodium
polystyrene sulfonate, sucrose, trataria caid,
tragacanth gum, triacetin, xanthan gum

Questions or comments? 1-800-632-6900

**DOSING**SHAKE WELL BEFORE USE.
Measure only with dosing cup provided.
Do not use dosing cup with other products.

Age (yr)	Dose
12 years	10 mL
to adult	EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use



## **PARENTS:** Learn about teen medicine abuse

www.StopMedicineAbuse.org

Do not use if carton is opened or printed foil under cap is broken or missing

COMPARE TO the active ingredient of DELSYM® ORANGE-FLAVOR \*See side panel NDC 30142-384-21



# Cough DM

Dextromethorphan Polistirex Extended-Release Oral Suspension

#### **COUGH SUPPRESSANT**

Contains Sodium Metabisulfite, a Sulfite That May Cause Allergic-Type Reactions

Day or Night Alcohol-Free Dosing Cup Included







38421 45 C5



ACTUAL SIZE



## Cough DM



-Flavored Liquid 3 FL OZ (89 mL)



dextromethorphan polistirex suspension

## **Product Information**

Product Type HUMAN OTC DRUG NDC:30142-384 Item Code (Source)

ORAL Route of Administration

## **Active Ingredient/Active Moiety**

**Ingredient Name Basis of Strength** Strength DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9 KYH) DEXTROMETHORPHAN 30 mg (DEXTROMETHORPHAN - UNII:7355X3ROTS) HYDROBROMIDE in 5 mL

Inactive Ingredients		
Ingredient Name	Strength	
POLISTIREX (UNII: 5H9 W9 GTW27)		
D&C RED NO. 30 (UNII: 2S42T2808B)		
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
GLYCERIN (UNII: PDC6A3C0OX)		
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)		
METHYLPARABEN (UNII: A2I8 C7HI9 T)		
POLYSORBATE 80 (UNII: 6OZP39ZG8H)		
POLYVINYL ACETATE (UNII: 32K497ZK2U)		
PO VIDONE, UNSPECIFIED (UNII: FZ989GH94E)		
PROPYLPARABEN (UNII: Z8IX2SC1OH)		
WATER (UNII: 059QF0KO0R)		
SODIUM METABISULFITE (UNII: 4VON5FNS3C)		
SODIUM POLYSTYRENE SULFONATE (UNII: 1699G8679Z)		
SUCRO SE (UNII: C151H8 M554)		
TARTARIC ACID (UNII: W48881119H)		
TRAGACANTH (UNII: 2944357O2O)		
TRIACETIN (UNII: XHX3C3X673)		
XANTHAN GUM (UNII: TTV12P4NEE)		

Product Characteristics		
Color	ORANGE	Score
Shape		Size
Flavor	ORANGE	Imprint Code
Contains		

	Packaging			
l	# Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date
l	1 NDC:30142-384-21	1 in 1 CARTON	08/28/2012	
	1	89 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091135	08/28/2012	

# Labeler - Kroger Company (006999528)

Revised: 5/2020 Kroger Company