

PERRIGO ALLERGY D- loratadine, pseudoephedrine sulfate tablet, film coated, extended release
L. Perrigo Company

Perrigo Allergy-D Drug Facts

Active ingredients (in each tablet)

Loratadine 5 mg

Pseudoephedrine sulfate 120 mg

Purpose

Antihistamine

Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - sneezing
 - itchy, watery eyes
 - runny nose
 - itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease

- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed.

Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not divide, crush, chew or dissolve the tablet

adults and children 12 years and over	1 tablet every 12 hours; not more than 2 tablets in 24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- **each tablet contains:** calcium 25 mg
- do not use if blister unit is broken or torn
- store between 20° to 25°C (68° to 77°F)
- keep in a dry place

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, pharmaceutical ink, povidone, titanium dioxide

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Claritin-D[®] 12 hour Extended Release Tablets

active ingredients

ALLERGY & CONGESTION

allergy-D

Pseudoephedrine Sulfate 120 mg/Nasal Decongestant

Loratadine 5 mg/Antihistamine

Extended-Release Tablets

12 Hour

INDOOR & OUTDOOR ALLERGIES

Relief of:

- Nasal and Sinus Congestion Due to Colds or Allergies
- Sneezing • Runny Nose • Itchy, Watery Eyes
- Itchy Throat or Nose Due to Allergies

Non-Drowsy*

*When taken as directed.

See Drug Facts Panel.

Actual Size

20 EXTENDED RELEASE TABLETS

Perrigo
 Allegan, MI 49010
 perigo.com

Distributed By

Drug Facts (continued)
 Questions or comments? 1-800-71-9-260



OPEN OTHER END

7U060 55 C3

CONVENIENT RE-CLOSING TAB

Drug Facts (continued)

Active ingredients (in each tablet)
 Loratadine 5 mg, Antihistamine
 Pseudoephedrine sulfate 120 mg, Nasal decongestant

Purpose

Uses
 temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 sneezing
 itchy, watery eyes
 runny nose
 itching of the nose or throat
 temporarily relieves nasal congestion due to a common cold, hay fever or other upper respiratory allergies
 reduces swelling of nasal passages
 temporarily relieves sinus congestion and pressure
 temporarily relieves throat breathing through the nose

Warnings
 Do not use
 if you have ever had an allergic reaction to this product or any of its ingredients
 if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, Parkinson's disease, or an MAOI) (certain drugs for depression, Parkinson's disease, or an MAOI), ask a doctor or pharmacist before taking this product.
 Ask a doctor before use if you have:
 heart disease
 thyroid disease
 high blood pressure
 trouble urinating due to an enlarged prostate gland
 diabetes
 liver or kidney disease
 Your doctor should determine if you need a different dose.

Other information
 Each tablet contains calcium 25 mg
 do not use if blister unit is broken or torn
 store between 20° to 25° C (68° to 77°)
 keep in a dry place

Inactive ingredients cross-cellulose sodium, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, pharmaceutical ink, povidone, titanium dioxide

Directions
 do not divide, crush, chew or dissolve the tablet
 adults and children
 1 tablet every 12 hours; not more than 2 tablets in 24 hours
 children under 12 years of age
 ask a doctor
 consumers with liver or kidney disease
 ask a doctor

Other information
 If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
 nervousness, dizziness or sleepiness occurs
 symptoms do not improve within 7 days or are accompanied by a fever
 an allergic reaction to the product occurs. Seek medical help right away.
 Stop use and ask a doctor if:
 Taking more than directed may cause drowsiness.
 When using this product do not take more than directed.

allergy-D 12 Hour

Compare to Claritin-D® 12 hour
 Extended Release Tablets
 active ingredients

NDC 0113-7777-60
 ALLERGY & CONGESTION

allergy-D

Pseudoephedrine Sulfate 120 mg/Nasal Decongestant
 Loratadine 5 mg/Antihistamine
 Extended-Release Tablets

INDOOR & OUTDOOR ALLERGIES 12 Hour

- Relief of:
- Nasal and Sinus Congestion Due to Colds or Allergies
 - Sneezing • Runny Nose • Itchy, Watery Eyes
 - Itchy Throat or Nose Due to Allergies

Non-Drowsy*
 *When taken as directed.
 See Drug Facts Panel

Actual Size



20 EXTENDED
 RELEASE TABLETS

PERRIGO ALLERGY D

loratadine, pseudoephedrine sulfate tablet, film coated, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0113-7777
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJ03BO7QN) (LORATADINE - UNII: 7AJ03BO7QN)	LORATADINE	5 mg
PSEUDOEPHEDRINE SULFATE (UNII: Y9DL7QPE6B) (PSEUDOEPHEDRINE - UNII: 7CUC9DDI9F)	PSEUDOEPHEDRINE SULFATE	120 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE (to off-white)	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	7U0
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-7777-60	20 in 1 CARTON	10/08/2021	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076050	10/08/2021	

