

TUKOL SUGAR FREE COUGH AND COLD- dextromethorphan hydrobromide and guaifenesin syrup
Genomma Lab USA

Tukol[®] Sugar Free Cough & Cold

Drug Facts

<i>Active ingredients (in each 10 mL)</i>	<i>Purposes</i>
Dextromethorphan HBr, 20 mg	Cough Suppressant
Guaifenesin, 200 mg	Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold,
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes.

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus),
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.

Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache.

These could be signs of a serious condition.

Do not exceed recommended dosage.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24 hour period,
- this adult strength product is not intended for use in children under 12 years of age,
- measure only with dosing cup provided,
- keep dosing cup with product,

- TSP = teaspoon
- mL = milliliter
- dose as follows:

age	dose
adults and children 12 years and over	10 mL (2 teaspoons) every 4 hours
children under 12 years	do not use

Other information

- store at 15-30°C (59-86°F). Do not refrigerate,
- alcohol-free,
- see bottom panel for lot number and expiration date.

Inactive ingredients

Anhydrous citric acid, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sucralose, xanthan gum.

Questions or comments?

1-877-994-3666 Monday to Friday from 8 am to 6 pm, Central Time.

Distributed by:
Genomma Lab USA Inc.,
Houston, TX 77098

PRINCIPAL DISPLAY PANEL - 118 mL Bottle Carton

Dextromethorphan HBr / Guaifenesin

Tukol[®]

SUGAR FREE

0%

COUGH & COLD

Ages

12+

HBP

HIGH BLOOD PRESSURE

RELIEVES COUGH

RELIEVES CHEST CONGESTION

SUGAR-FREE FORMULA

FOR DIABETICS

4 FL OZ (118 mL)

Tukol[®]

SUGAR FREE
SIN AZÚCAR

**DO NOT USE IF PRINTED SEAL UNDER CAP
IS TORN OR MISSING**
**NO SE USE SI EL SELLO IMPRESO DEBAJO DE LA TAPA
ESTA ROTO O FALTA**

Dextromethorphan HBr / Guaifenesin
Dextrometorfano HBr / Guaifenesina

Tukol[®]

SUGAR FREE
SIN AZÚCAR **0%**

COUGH & COLD
TOS Y RESFRIADO

Age
12+



RELIEVES COUGH /
ALIVIA TOS

RELIEVES CHEST CONGESTION /
ALIVIA CONGESTIÓN DE PECHO

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, 20 mg..... Cough Suppressant
Guaifenesin, 200 mg..... Expectorant

Purposes

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold,
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes.

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus),
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.

Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache.

These could be signs of a serious condition.

Do not exceed recommended dosage.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24 hour period,
- this adult strength product is not intended for use in children under 12 years of age,
- measure only with dosing cup provided,
- keep dosing cup with product.

**SUGAR-FREE FORMULA
FOR DIABETICS**
FÓRMULA "CERO AZÚCAR" PARA DIABÉTICOS

4 FL OZ (118 mL)

■ TSP = teaspoon ■ mL = milliliter
■ dose as follows:

age	dose
adults and children 12 years and over	10 mL (2 teaspoons) every 4 hours
children under 12 years	do not use

Drug Facts (continued)

Other information

- store at 15-30°C (59-86°F). Do not refrigerate,
- alcohol-free,
- see bottom panel for lot number and expiration date.

Inactive ingredients

Anhydrous citric acid, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sucralose, xanthan gum.

Questions or comments?

1-877-994-3666 Monday to Friday from 8 am to 6 pm, Central Time.

Información del Medicamento

**Ingredientes activos Propósitos
(en cada 10 mL)**

Dextrometorfano HBr, 20 mg.....Supresor de la Tos
Guaifenesina, 200 mgExpectorante

Usos

- alivia temporalmente la tos debida a una irritación menor de garganta y bronquial que puede ocurrir con un resfriado,
- ayuda a aflojar la flema (mucosidad) y a adelgazar las secreciones bronquiales para drenar los conductos bronquiales.

**Información del Medicamento
(continuación)**

Suspenda su uso y consulte a su médico si la tos dura más de 7 días, vuelve, o está acompañada de fiebre, dolor de cabeza persistente o erupción cutánea. Estos podrían ser signos de una condición grave.

No exceder la dosis recomendada.

Si está embarazada o amamantando, consulte a un profesional de la salud antes de usar.

Manténgase fuera del alcance de los niños. En caso de sobredosis, solicite ayuda médica o póngase en contacto inmediatamente con un Centro de Control Envenenamiento (1-800-222-1222).

Indicaciones

- no tomar más de 6 dosis en un período de 24 horas,
- este producto de concentración para adultos no está previsto para su uso en niños menores de 12 años de edad,
- medir solo con la copa dosificadora incluida,
- mantenga la copa dosificadora con este producto,
- TSP = cucharadita ■ mL = mililitro,
- administrar dosis según se indica a continuación:

edad	dosis
adultos y niños 12 años de edad y mayores	10 mL (2 cucharaditas) cada 4 horas
niños menores de 12 años	no utilizar

Información adicional

- almacenar entre 15-30 °C (59-86 °F). No refrigerar.
- libre de alcohol.
- lea el panel del fondo para el Lote y Fecha de Expiración.

Advertencias

No lo use si está tomando un medicamento de prescripción inhibidor de la monoamino oxidasa (MAO) (ciertos medicamentos para la depresión, condiciones psiquiátricas, emocionales o para la enfermedad de Parkinson) o las 2 semanas después de suspender medicamento MAO. Si no sabe si su medicamento contiene un MAO, pregunte a un médico o farmacéutico antes de tomar este producto.

Pregunte a un doctor antes de usar si tiene

- tos que se produce con mucha flema (mucosidad),
- tos duradera o crónica, tal como ocurre al fumar, asma, bronquitis crónica o enfisema.

Ingredientes inactivos

Acido cítrico anhidro, saborizantes, glicerina, propilenglicol, agua purificada, benzoato de sodio, sucralosa, goma xantana.

¿Preguntas o comentarios?

1-877-994-3666 Lunes a Viernes de 8 am a 6 pm, Tiempo del Centro.

Distributed by / Distribuido por:
Genomma Lab USA Inc.,
Houston, TX 77098



2000012980

Rev. 1



EXP:

LOT No./LOTE:

TUKOL SUGAR FREE COUGH AND COLD

dextromethorphan hydrobromide and guaifenesin syrup

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50066-505
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	2 mg in 1 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	20 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50066-505-04	1 in 1 CARTON	06/13/2024	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/13/2024	

Labeler - Genomma Lab USA (832323534)

Revised: 10/2024

Genomma Lab USA