

**MUCUS RELIEF COLD AND SINUS MAXIMUM STRENGTH- acetaminophen,  
guaifenesin, phenylephrine hcl liquid  
MEIJER, INC.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredients (in each 20 mL)**

**Acetaminophen 650 mg**

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

**Purposes**

Pain reliever/fever reducer

Expectorant

Nasal decongestant

**Uses**

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - sore throat
  - headache
  - nasal congestion
  - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occur with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

**When using this product,**

**do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or last more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or persistent headache. These could be a signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL=milliliter
- adults and children 12 years and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

### **Other information**

- each 20 mL contains: **sodium 12 mg**
- store at 20-25°C (68-77°F). Do not refrigerate

### **Inactive ingredients**

anhydrous citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum

### **Questions or comments?**

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

### **Principal Display Panel**

Compare to the active ingredient in Mucinex® Fast-Max® Maximum Strength Cold & Sinus Liquid active ingredients\*

Cold & Sinus

#### **Mucus Relief**

Aches & Fever (Acetaminophen)

Chest Congestion & Mucus (Guaifenesin)

Stuffy Nose (Phenylephrine HCl)

Maximum Strength

Relieves:

Aches & Fever

Chest Congestion

Mucus

Stuffy Nose

For Ages 12 & Over

FL OZ (mL)

DIST.BY MEIJER DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

[www.meijer.com](http://www.meijer.com)

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex® Fast Max® Maximum Strength Cold & Sinus Liquid.

**Package Label**

NDC 41250-136-06



Compare to Mucinex® Fast-Max® Maximum Strength Cold & Sinus Liquid active ingredients\*

Cold & Sinus

# Mucus Relief

Aches & Fever (Acetaminophen)  
Chest Congestion & Mucus (Guaifenesin)  
Stuffy Nose (Phenylephrine HCl)

Maximum Strength

Relieves:  
Aches & Fever  
Chest Congestion  
Mucus  
Stuffy Nose

6 FL OZ  
(177 mL)

• For ages 12 & over

PLD-B281B LB002913

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex® Fast-Max® Maximum Strength Cold & Sinus Liquid.

PLD-B281B  
LB002912



DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com

### Drug Facts

#### Active ingredients (in each 20 mL)

Acetaminophen 650 mg.....Pain reliever/fever reducer  
Guaifenesin 400 mg.....Expectorant  
Phenylephrine HCl 10 mg.....Nasal decongestant

**Uses** ■ temporarily relieves these common cold and flu symptoms: ■ minor aches and pains  
■ sore throat ■ headache ■ nasal congestion  
■ sinus congestion and pressure  
■ temporarily reduces fever  
■ helps loosen phlegm (mucus) and thin bronchial

PEEL CORNER FOR MORE DRUG FACTS

### Drug Facts (continued)

secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### Warnings

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### Drug Facts (continued)

- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- liver disease ■ heart disease ■ diabetes
- thyroid disease ■ high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

PEEL CORNER FOR MORE DRUG FACTS

### Drug Facts (continued)

- cough comes back, or occurs with rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- adults and children 12 years and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

### Drug Facts (continued)

#### Other information

- each 20 mL contains: sodium 12 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** anhydrous citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum

**Questions or comments?** Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Meijer Mucus Relief Cold & Sinus Maximum Strength Liquid

**MUCUS RELIEF COLD AND SINUS MAXIMUM STRENGTH**

acetaminophen, guaifenesin, phenylephrine hcl liquid

## Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:41250-136 |
| <b>Route of Administration</b> | ORAL           |                           |               |

## Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength              | Strength           |
|---|--------------------------------|--------------------|
| <b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)               | ACETAMINOPHEN                  | 650 mg<br>in 20 mL |
| <b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)                   | GUAIFENESIN                    | 400 mg<br>in 20 mL |
| <b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE<br>HYDROCHLORIDE | 10 mg<br>in 20 mL  |

## Inactive Ingredients

| Ingredient Name                                       | Strength |
|---|----------|
| <b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)       |          |
| <b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)            |          |
| <b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)         |          |
| <b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)         |          |
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)                    |          |
| <b>PROPYL GALLATE</b> (UNII: 8D45NN7V92)              |          |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)            |          |
| <b>WATER</b> (UNII: 059QF0KO0R)                       |          |
| <b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)             |          |
| <b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K) |          |
| <b>SORBITOL</b> (UNII: 506T60A25R)                    |          |
| <b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)                   |          |
| <b>XANTHAN GUM</b> (UNII: TTV12P4NEE)                 |          |

## Packaging

| # | Item Code        | Package Description  | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:41250-136-06 | 177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 04/30/2015           | 12/31/2024         |

## Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341                                  | 04/30/2015           | 12/31/2024         |

**Labeler** - MEIJER, INC. (006959555)

