

CRANE SAFETY ASPIRIN- aspirin tablet, coated
CRANE SAFETY ASPIRIN- aspirin tablet, film coated
THOMPSON ASPIRIN- aspirin tablet, film coated
Thompson

Thompson Aspirin

Drug Facts

Active ingredient

Aspirin (NSAID*) 325 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- if ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose

Adults and children: (12 years and older)

Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours.

Children under 12 years:

Consult a doctor

Other information

- store at room temperature

- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments?

877.506.4291

Crane Safety Aspirin Label

5 Grain (325 mg)

ASPIRIN

PainReliever/Fever Reducer (NSAID)

CRANE Safety®

Pull To Open

This Package Is For Households Without Young Children.

Aches, Fever

Aspirin (NSAID) 325 mg

100 Tablets

Tamper Evident

Sealed Packets



Crane Safet Aspirin label

5 Grain (325 mg)

ASPIRIN

PainReliever/Fever Reducer (NSAID)

CRANE Safety®

Pull To Open

This Package Is For Households Without Young Children.

Aches, Fever

Aspirin (NSAID) 325 mg

100 Tablets

Tamper Evident

Sealed Packets

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purpose Aspirin 325 mg (NSAID) Pain reliever/fever reducer nonsteroidal anti-inflammatory drug</p> <p>Uses Temporarily relieves minor aches and pains associated with: ■ headache ■ muscle pain ■ minor pain of arthritis ■ colds ■ toothache ■ menstrual pain Temporarily reduces fever.</p> <p>Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock If an allergic reaction occurs, stop use and seek medical help right away.</p>	<p>Drug Facts (continued)</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcohol drinks every day while using this product ■ take more or for a longer time than directed</p> <p>Do not use ■ if you are allergic to aspirin or to any other pain reliever/fever reducer ■ if you ever had an allergic reaction to this product or any of its ingredients</p> <p>Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn, ulcers, or kidney disease ■ you are taking a diuretic ■ you have asthma (Continued on opposite panel)</p>	<p>5 GRAIN (325 MG) ASPIRIN</p> <p>PAIN RELIEVER/FEVER REDUCER (NSAID)</p> <p>CRANE Safety</p> <p>Pull To Open</p> <p>THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.</p> <p>ACHES, FEVER ASPIRIN (NSAID) 325 MG</p> <p>100 Tablets TAMPER EVIDENT SEALED PACKETS</p>	<p>Drug Facts (continued)</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis</p> <p>Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ if ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area ■ any new symptoms appear</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p>Drug Facts (continued)</p> <p>Directions ■ drink a full glass of water with each dose</p> <p>Adults and children (12 years and older) Take 1 or 2 tablets with water every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours.</p> <p>Children under 12 years: Do not give to children under 12 years of age unless directed by a doctor.</p> <p>Other information ■ store at 77°F (25°C); excursions permitted between 59-86°F (15-30°C) ■ avoid excessive heat and humidity ■ tamper evident sealed packets ■ do not use any opened or torn packets ■ use by expiration date on package</p> <p>Inactive ingredients corn starch, croscarmellose sodium*, hypromellose, microcrystalline cellulose*, mineral oil*, polyethylene glycol*, propylene glycol*, titanium dioxide* may contain*</p> <p>Questions or comments? 877.508.4291</p>	<p>5 GRAIN (325 MG) ASPIRIN</p> <p>PAIN RELIEVER/FEVER REDUCER (NSAID)</p> <p>CRANE Safety</p> <p>Reorder C590533</p> <p>Distributed by Crane Safety Houston, TX 77073 877.508.4291 www.cranesafetyco.com</p> <p>0 73408 90533 8</p>
---	--	---	---	--	---

5 Grain (325 mg)

ASPIRIN

PainReliever/Fever Reducer (NSAID)

CRANE Safety®

Pull To Open

This Package Is For Households Without Young Children.

Aches, Fever

Aspirin (NSAID) 325 mg

100 Tablets

Tamper Evident

Sealed Packets

5 GRAIN (325 MG) ASPIRIN

PAIN RELIEVER/FEVER REDUCER (NSAID)

Pull to
Open

ACHES, FEVER
ASPIRIN (NSAID) 325 MG

THIS PACKAGE IS FOR HOUSEHOLDS
WITHOUT YOUNG CHILDREN

100 TABLETS

TAMPER EVIDENT
SEALED PACKETS

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin 325 mg (NSAID) Pain reliever/fever reducer
nonsteroidal anti-inflammatory drug

Uses
temporarily relieves
■ headache ■ muscle pain ■ minor pain of arthritis
■ pain and fever of colds ■ toothache ■ menstrual pain

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcohol drinks every day while using this product
■ take more or for a longer time than directed

Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you are taking a diuretic
■ you have asthma

Drug Facts (continued)
Ask a doctor or pharmacist before use if you are taking a prescription drug for:
■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood
 ■ have bloody or black stools
■ pain gets worse or lasts more than 10 days
■ fever lasts more than 3 days
■ ringing in the ears or loss of hearing occurs
■ redness or swelling is present
■ new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ drink a full glass of water with each dose

Adults and children (12 years and older)	Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
Children under 12 years:	Consult a doctor

Other information
■ store at room temperature
■ avoid excessive heat and humidity
■ tamper evident sealed packets
■ do not use any opened or torn packets

Inactive ingredients
corn starch, hypromellose*, microcrystalline cellulose*, polyethylene glycol*, povidone*, propylene glycol*
*may contain

Questions or comments? 877.506.4291

5 GRAIN (325 MG) ASPIRIN

PAIN RELIEVER/FEVER REDUCER (NSAID)

100 TABLETS

TAMPER EVIDENT
SEALED PACKETS

Retain carton for complete product information

5 Grain (325 mg)
ASPIRIN
Thompson
Safety Works Business Thrives™
Pull To Open
This package is for households without young children.
Aches, Fever
Aspirin (NSAID) 325 mg
100 Tablets
Tamper Evident
Sealed Packets

5 GRAIN (325 MG) ASPIRIN

PAIN RELIEVER/FEVER REDUCER (NSAID)

Pull to
Open

THIS PACKAGE IS FOR HOUSEHOLDS
WITHOUT YOUNG CHILDREN.

ACHES, FEVER
ASPIRIN (NSAID) 325 MG

100 TABLETS

TAMPER EVIDENT
SEALED PACKETS

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin 325 mg (NSAID) Pain reliever/fever reducer
nonsteroidal anti-inflammatory drug

Uses
temporarily relieves
■ headache ■ muscle pain ■ minor pain of arthritis
■ pain and fever of colds ■ toothache ■ menstrual pain

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Drug Facts (continued)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcohol drinks every day while using this product
■ take more or for a longer time than directed

Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you are taking a diuretic
■ you have asthma

Drug Facts (continued)
Ask a doctor or pharmacist before use if you are taking a prescription drug for:
■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood
 ■ have bloody or black stools
■ pain gets worse or lasts more than 10 days
■ fever lasts more than 3 days
■ ringing in the ears or loss of hearing occurs
■ redness or swelling is present
■ new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Directions
■ drink a full glass of water with each dose

Adults and children (12 years and over)	Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
Children under 12 years:	Consult a doctor

Other information
■ store at room temperature
■ avoid excessive heat and humidity
■ tamper evident sealed packets
■ do not use any opened or torn packets

Inactive ingredients
corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments? 877.506.4291

5 GRAIN (325 MG) ASPIRIN

PAIN RELIEVER/FEVER REDUCER (NSAID)

100 TABLETS

TAMPER EVIDENT
SEALED PACKETS

Retain carton for complete product information

CRANE SAFETY ASPIRIN

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73408-620	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	325 mg	
Inactive Ingredients				
Ingredient Name			Strength	
STARCH, CORN (UNII: O8232NY3SJ)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POVIDONE (UNII: FZ989GH94E)				
Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	FR21	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-620-33	50 in 1 BOX	02/10/2021	04/01/2024
1		2 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	02/10/2021	04/01/2024	

CRANE SAFETY ASPIRIN				
aspirin tablet, film coated				
Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73408-099	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients	
Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
STARCH, CORN (UNII: O8232NY3SJ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	TCL;011
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-099-33	50 in 1 BOX	01/17/2020	08/31/2022
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/17/2020	08/31/2022

THOMPSON ASPIRIN

aspirin tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73408-618
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white	Score	no score
Shape	RECTANGLE	Size	10mm
Flavor		Imprint Code	Aspirin;44;157
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-618-33	50 in 1 BOX	05/15/2024	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/15/2024	

CRANE SAFETY ASPIRIN

aspirin tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73408-616
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)

STARCH, CORN (UNII: O8232NY3SJ)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	Aspirin;44;157
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-616-33	50 in 1 BOX	01/17/2020	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/17/2020	08/31/2024

Labeler - Thompson (080998015)

Registrant - Unifirst First Aid Corporation (832947092)

Establishment

Name	Address	ID/FEI	Business Operations
Time Cap Laboratories, Inc.		037052099	manufacture(73408-099)

Establishment

Name	Address	ID/FEI	Business Operations
Medique Products		086911794	pack(73408-618)

Revised: 9/2025

Thompson