

**ENTERIC COATED ASPIRIN REGULAR STRENGTH- aspirin tablet, delayed release**  
**Safeway**

-----

**SIG - 1081B - 2024-1125**

***Drug Facts***

**Active ingredient (in each tablet)**

Aspirin 325 mg (NSAID <sup>1</sup>)

---

1 nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

- for the temporary relief of minor aches and pains due to:
  - headache
  - muscle pain
  - toothache
  - menstrual pain
  - colds
  - minor pain of arthritis
- or as recommended by your doctor

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for:

- diabetes
- gout
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs

These could be signs of a serious condition

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- drink a full glass of water with each dose

- adults and children 12 years and over: take 1 to 2 tablets every 4 hours; do not exceed 12 tablets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information

**Inactive ingredients**

corn starch, croscarmellose sodium, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

**PRINCIPAL DISPLAY PANEL**

COMPARE TO

Ecotrin® Regular Strength Aspirin

Active ingredient†

NDC 21130-081-05

Signature SELECT™

REGULAR STRENGTH

ENTERIC COATED

ASPIRIN 325mg

Pain Reliever (NSAID)

- Safety coated

Actual Size

500 ENTERIC COATED TABLETS

**COMPARE TO**  
Ecoirin® Regular Strength Aspirin  
Active Ingredient\*

NDC 21130-081-05

**Signature SELECT™**

**REGULAR STRENGTH  
ENTERIC COATED  
ASPIRIN 325mg**

Pain Reliever (NSAID)  
• Safety coated

Actual Size 

**500 ENTERIC  
COATED TABLETS**

**DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts**  
Aspirin 325mg (NSAID)\*, nonsteroidal anti-inflammatory drug  
**Active ingredient (in each tablet)**.....Pain reliever

**Uses**  
■ for the temporary relief of minor aches and pains due to:  
■ muscle pain ■ toothache ■ menstrual pain ■ headache  
■ or as recommended by your doctor ■ colds ■ minor pain of arthritis

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:  
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed

**Do not use if you are allergic to aspirin or any other pain reliever/fever reducer**

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you ■ you are taking a diuretic  
■ you have a history of stomach problems, such as heartburn ■ you have asthma  
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

**Ask a doctor or pharmacist before use if you are taking a prescription drug for:**  
■ diabetes ■ gout ■ arthritis

**Stop use and ask a doctor if**  
■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood  
■ have bloody or black stools ■ have stomach pain that does not get better  
■ pain gets worse or lasts for more than 10 days ■ redness or swelling is present  
■ fever gets worse or lasts for more than 3 days ■ new symptoms occur  
■ ringing in the ears or loss of hearing occurs  
These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**  
■ drink a full glass of water with each dose  
■ adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last; do not take more than 12 tablets in 24 hours unless directed by a doctor  
■ children under 12 years: ask a doctor

**Other information**  
■ store between 20°-25°C (68°-77°F) in a dry place

**Inactive ingredients**  
corn starch, croscarmellose sodium, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polyorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

**DISTRIBUTED BY:**  
BETTER LIVING BRANDS LLC  
P.O. BOX 99  
PLEASANTON, CA 94666-0099  
1-888-725-3829

**Scan for info**  
RD 24215  
S4972  
21130178261 2  
1-800-835-5630

# ENTERIC COATED ASPIRIN REGULAR STRENGTH

aspirin tablet, delayed release

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:21130-081
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

## Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	T
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-081-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	08/05/2015	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/05/2015	

**Labeler** - Safeway (009137209)

Revised: 11/2024

Safeway