

**SALICYLIC ACID- liquid corn and callus remover liquid**  
**Foodhold USA**

-----  
**CareOne Liquid Corn & Callus Remover**

**Active ingredient**

Salicylic acid 17%w/w

**Purpose**

Corn and callus remover

**Uses**

- for the removal of corns and calluses
- relieves pain by removing corns and calluses

**Warnings**

**For external use only.**

**Flammable**

- keep away from fire or flame
- cap bottle tightly and store at room temperature away heat

**Do not use**

- on irritated skin
- on any area that is infected or reddened
- if you are a diabetic, or if you have poor blood circulation, except under the advice and supervision of a doctor or podiatrist

**When using this product**

- avoid contact with eyes
- avoid inhaling vapors
- if product gets into the eyes, flush with water for 15 minutes

**Stop and ask a doctor if**

discomfort persists, see your doctor or podiatrist

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- wash affected area and dry thoroughly

- apply one drop at a time with applicator, to sufficiently cover each corn/callus; let dry
- use cushions with hole centered over corn or callus to relieve pressure (additional cushions may be purchased separately)
- repeat this procedure once or twice daily as needed for up to 14 days (until corn/callus is removed)

### **Other information**

store between 20°C to 30°C (68°F to 86°F)

### **Inactive ingredients**

camphor, castor oil, ethanol, ethyl ether, nitrocellulose

### **Questions?**

Call 1-888-423-0139

### **Principal Display Panel**

**CAREONE**

**LIQUID CORN & CALLUS REMOVER**

Salicylic Acid

REMOVES CORNS AND CALLUSES SAFELY

SOFT FOAM CUSHIONS AND PROTECTS

**NET WT 0.31 FL OZ (9.8 mL)**

**3 CUSHIONS**

CAREone™

Compare to the Active Ingredient in Dr. Scholl's Liquid Corn/Callus Remover\*

# LIQUID CORN & CALLUS REMOVER

Salicylic Acid

REMOVES CORNS AND CALLUSES SAFELY  
SOFT FOAM CUSHIONS AND PROTECTS

OUR PHARMACISTS RECOMMEND

NET WT. 0.31 FL OZ (9.8 mL) 3 CUSHIONS

CAREone™

## LIQUID CORN & CALLUS REMOVER

**Drug Facts**

Active ingredient	Purpose
Salicylic acid 17% w/w	Corn and callus remover

**Uses**

- for the removal of corns and calluses
- relieves pain by removing corns and calluses

**Warnings**

For external use only.

**Flammable**

- keep away from fire or flame
- cap bottle tightly and store at room temperature away from heat

**Do not use**

- on irritated skin
- on any area that is infected or reddened
- if you are a diabetic, or if you have poor blood circulation, except under the advice and supervision of a doctor or podiatrist

**When using this product**

- avoid contact with eyes
- avoid inhaling vapors
- if product gets into the eye, flush with water for 15 minutes

**Stop use and ask a doctor if discomfort persists, see your doctor or podiatrist.**

**Keep out of reach of children.**  
If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- wash the affected area and dry thoroughly
- apply one drop at a time with applicator to sufficiently cover each corn/callus; let dry
- use cushion with hole centered over corn or callus to relieve pressure (additional cushions may be purchased separately)
- repeat this procedure once or twice daily as needed for up to 14 days (until corn/callus is removed)

**Other information**

- store between 20° - 25°C (68° - 77°F)
- close tightly to prevent solidification
- store in a cool place

**Inactive ingredients** camphor, castor oil, ethanol, ethyl ether, nitrocellulose

**Questions?** Call 1-866-964-0939

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Dr. Scholl's®.

DISTRIBUTED BY FOODHOLD U.S.A., LLC  
LANDOVER, MD 20785 1-877-846-9949  
© 2017 Ahold Licensing Sarl  
MADE IN USA WITH U.S. AND IMPORTED PARTS

Quality guaranteed or your money back.

41-030CO-01

CARE: WIPE TOP OF BOTTLE  
CLEAN AFTER EACH USE

FPO

0 41520 04533 1

IMPORTANT: CLOSE CAP TIGHTLY AFTER USE

NDC 00000-000-00

CAREone™

# LIQUID CORN & CALLUS REMOVER

SALICYLIC ACID  
REMOVES CORNS AND CALLUSES SAFELY  
AND EFFECTIVELY

NET WT.  
0.31 FL OZ (9.8 mL)

**Active ingredient:** Salicylic acid (17% w/w)  
**Purpose:** Corn and callus remover  
**Uses:** For the removal of corns and calluses. Relieves pain by removing corns and calluses.  
**Warnings:** For external use only. **Flammable:** Keep away from fire or flame. **Cap bottle tightly** and store at room temperature away from heat. **Do not use** this product if you are diabetic or have poor blood circulation, except under the advice and supervision of a doctor or podiatrist.

**Do not use this product** on irritated skin, on any area that is infected or reddened. **When using this product** avoid contact with eyes, if product gets into the eyes, flush with water for 15 minutes. Avoid inhaling vapors. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away. **Directions:** Wash affected area and dry thoroughly. Apply one drop at a time with applicator to sufficiently cover each corn/callus. Let dry. Use cushions with hole centered over corn or callus to relieve pressure. Repeat this procedure once or twice daily as needed for up to 14 days (until corn/callus is removed). **Other information:** Store at 20° - 25° C (68° - 77° F). Close tightly to prevent solidification. Store in a cool place. **Inactive ingredients:** camphor, castor oil, ethanol, ethyl ether, nitrocellulose **Questions?** Call 1-866-964-0939

**Keep packaging for complete drug facts information**  
**MADE IN USA WITH U.S. AND IMPORTED PARTS**  
DISTRIBUTED BY FOODHOLD U.S.A., LLC  
LANDOVER, MD, 20785 1-877-846-9949  
© 2017 Ahold Licensing Sarl

QUALITY GUARANTEED  
OR YOUR MONEY BACK.

50-030CO

## SALICYLIC ACID

liquid corn and callus remover liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41520-569
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	0.17 mg in 9.8 g

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>PYROXYLIN</b> (UNII: KYR8BR2X6O)	
<b>CAMPHOR (NATURAL)</b> (UNII: N20HL7Q941)	
<b>CASTOR OIL</b> (UNII: D5340Y2I9G)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:41520-569-03	9.8 g in 1 PACKAGE; Type 0: Not a Combination Product	10/20/2017	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M030	10/20/2017	

**Labeler** - Foodhold USA (809183973)

Revised: 1/2024

Foodhold USA