

**DIMETAPP NIGHTTIME COLD AND CONGESTION- diphenhydramine hydrochloride and phenylephrine hydrochloride solution**  
**Foundation Consumer Brands**

-----  
**Dimetapp®**  
**Nighttime Cold and Congestion**

**Drug Facts**

<b>Active ingredients (in each 10 mL)</b>	<b>Purposes</b>
Diphenhydramine HCl, 12.5 mg	Antihistamine/cough suppressant
Phenylephrine HCl, 5 mg	Nasal decongestant

**Uses**

- temporarily relieves:
  - nasal congestion
  - runny nose
  - cough
  - sneezing
  - itching of the nose or throat
  - itchy, watery eyes due to hay fever
- temporarily restores freer breathing through the nose

**Warnings**

**Do not use**

- to sedate a child or to make a child sleepy
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other product containing diphenhydramine, even one used on skin

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are**

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

**When using this product**

- **do not use more than directed**
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occurs
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

<b>age</b>	<b>dose</b>
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

**Other information**

- each 10 mL contains: **sodium 8 mg**
- store at 20-25°C (68-77°F)
- do not refrigerate

**Inactive ingredients**

anhydrous citric acid, artificial flavor, FD&C blue no.1, FD&C red no. 40, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

## Questions or comments?

Call **1-888-594-0828** weekdays 9 AM to 5 PM EST

Distributed by:  
Foundation Consumer Brands, LLC  
Pittsburgh, PA 15212

## **PRINCIPAL DISPLAY PANEL - 118 mL Bottle Carton**

CHILDREN'S  
Dimetapp®

DIPHENHYDRAMINE HCl (Antihistamine/Cough Suppressant)  
PHENYLEPHRINE HCl (Nasal Decongestant)

PHARMACIST  
RECOMMENDED

Nighttime  
Cold &  
Congestion

Relieves + comforts:

- ✓ Stuffy, runny nose
- ✓ Itchy, watery eyes
- ✓ Sneezing
- ✓ Cough

6+  
YRS

4 FL OZ  
(118 mL)

Grape Flavor • Alcohol Free



## DIMETAPP NIGHTTIME COLD AND CONGESTION

diphenhydramine hydrochloride and phenylephrine hydrochloride solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:80070-340
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40)	DIPHENHYDRAMINE	6.25 mg

(DIPHENHYDRAMINE - UNII:8GTS82S83M)	HYDROCHLORIDE	in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYL GALLATE</b> (UNII: 8D4SNN7V92)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Product Characteristics

<b>Color</b>	PURPLE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:80070-340-04	1 in 1 CARTON	09/15/2021	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M012	09/15/2021	

**Labeler** - Foundation Consumer Brands (117603632)