

**BAEKCHO-DS- licorice ext syrup**  
**I World Pharmaceutical Co., Ltd.**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).*

-----

**Drug Facts**

Licorice Ext

Glycerine,menthol oil,sucralose,xylitol,purified water,etc

(Loss of appetite), stomach bloating, dyspepsia, overeating, nausea, vomiting, suicide, constipation, dilute stools,

keep out of reach of the children

Adult and 15 years and older: 1 time 10 mL

Pediatric patients were divided into two groups: 1 to 11 years old, 6.5 mL, 8 to 10 years old, 5.0 mL, 5 to 7 years old, 3.5 mL, 3-4 years old, 2.5 mL,

Take it three times a day after meals. The interval should be 4 hours or more.

1. Do not take this medicine.

Do not take to infants (infants, babies) under 3 months. In addition, infants (infants and young babies) who are under the age of one for at least three months must give priority to receiving medical attention, and should not be taken unless they are inevitable.

2. Consult a doctor, dentist or pharmacist before taking this medicine.

1) patients with hypertension, elderly (elderly), cardiac or renal (kidney) patients, edema (swelling) patients

2) People who are taking other medicines: potassium-containing preparations, licorice-containing preparations, glycyrrhizic acid or its salt-containing preparations, loop-type diuretics (furosemide, ethacrynic acid) or thiazide diuretics (trichloromethiazide) (Combined with) Aldosteronism or hypokalemia due to myocarditis (myopathy) is likely to occur due to taking it carefully.

3) Patients receiving medical treatment

3. Immediately stop taking this drug and consult a doctor, dentist or pharmacist if: Have this attached document at the time of consultation.

1) Upper aldosteronism: If you have decreased urine volume, face and hands and feet are swollen, your eyelids are getting heavy, your hands are hardening, your blood pressure is getting high or you have a headache (such as a maximum dose of 1g per day as a licorice, Observation (measurement of serum potassium levels) should be sufficient and should be discontinued if abnormality is confirmed, because gastric aldosteronism such as elevation of blood pressure, elevation of blood pressure, retention of sodium body fluid (swelling), swelling (swelling)

2) Myopathy (myopathy): As a result of hypokalemia, myopathy (myopathy) may occur. If the observation is sufficient and anomalies such as helplessness, cramps and paralysis are confirmed

3) If there is no improvement in symptoms even for several days

4. Other cautions when taking this medicine

1) Keep the prescribed dosage and dosage.

2) When taking a child, take it under the supervision of a guardian.

3) Do not take it for a long time.

5. Storage Precautions

1) Keep out of reach of children.

2) Taking medicines out of their original containers and storing them in other containers may cause accidents or poor quality of medicines due to misuse (misuse), so keep them in their original container

and keep them tightly closed.

3) Avoid direct sunlight and store in a cool, dry place.

for oral use only

## BAEKCHO-DS SYRUP

<b>Drug Facts</b>	
<b>Active Ingredients (in 1ml)</b>	<b>Purpose</b>
Licorice ext 10mg -----	
<b>Uses</b> (Loss of appetite), stomach bloating, dyspepsia, overeating, nausea, vomiting, suicide, constipation, dilute stools,	
<b>Warnings</b> <b>1. Do not take this medicine.</b> Do not take to infants (infants, babies) under 3 months. In addition, infants (infants and young babies) who are under the age of one for at least three months must give priority to receiving medical attention, and should not be taken unless they are inevitable. <b>2. Consult a doctor, dentist or pharmacist before taking this medicine.</b> 1) patients with hypertension, elderly (elderly), cardiac or renal (kidney) patients, edema (swelling) patients 2) People who are taking other medicines: potassium-containing preparations, licorice-containing preparations, glycyrrhizic acid or its salt-containing preparations, loop-type diuretics (furosemide, ethacrynic acid) or thiazide diuretics (trichloromethiazide) (Combined with) Aldosteronism or hypokalemia due to myocarditis (myopathy) is likely to occur due to taking it carefully. 3) Patients receiving medical treatment <b>3. Immediately stop taking this drug and consult a doctor, dentist or pharmacist if: Have this attached document at the time of consultation.</b> 1) Upper aldosteronism: If you have decreased urine volume, face and hands and feet are swollen, your eyelids are getting heavy, your hands are hardening, your blood pressure is getting high or you have a headache (such as a maximum dose of 1g per day as a licorice, Observation (measurement of serum potassium levels) should be sufficient and should be discontinued if abnormality is confirmed, because gastric aldosteronism such as elevation of blood pressure, elevation of blood pressure, retention of sodium body fluid (swelling), swelling (swelling) 2) Myopathy (myopathy): As a result of hypokalemia, myopathy (myopathy) may occur. If the observation is sufficient and anomalies such as helplessness, cramps and paralysis are confirmed 3) If there is no improvement in symptoms even for several days <b>4. Other cautions when taking this medicine</b> 1) Keep the prescribed dosage and dosage. 2) When taking a child, take it under the supervision of a guardian. 3) Do not take it for a long time. <b>5. Storage Precautions</b> 1) Keep out of reach of children. 2) Taking medicines out of their original containers and storing them in other containers may cause accidents or poor quality of medicines due to misuse (misuse), so keep them in their original container and keep them tightly closed. 3) Avoid direct sunlight and store in a cool, dry place.	
<b>Directions</b> ■ Adult and 15 years and older: 1 time 10 ml. ■ Pediatric patients were divided into two groups: 1 to 11 years old, 6.5 ml, 8 to 10 years old, 5.0 ml, 5 to 7 years old, 3.5 ml, 3-4 years old, 2.5 ml ■ Take it three times a day after meals. The interval should be 4 hours or more	
<b>Other Information</b> ■ Store at controlled room temperature 20-25 degrees C (68-77 degrees F).	
<b>Inactive Ingredient</b> Glycerine, menthol, oil, sucralose, xylitol, purified water, etc	

## BAEKCHO-DS

licorice ext syrup

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:73442-0007	
<b>Route of Administration</b>	ORAL			
<b>Active Ingredient/Active Moiety</b>				
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>	
LICORICE (UNII: 61ZBX54883) (LICORICE - UNII:61ZBX54883)		LICORICE	10 g in 1 mL	
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>			<b>Strength</b>	
GLYCERIN (UNII: PDC6A3C0OX)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
MENTHOL (UNII: L7T10EIP3A)				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:73442-0007-1	100 mL in 1 BOTTLE, GLASS; Type 0: Not a Combination Product	03/29/2019	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
unapproved drug other		03/23/2019		

**Labeler** - IWorld Pharmaceutical Co., Ltd. (688222857)

**Registrant** - IWorld Pharmaceutical Co., Ltd. (688222857)

### Establishment

Name	Address	ID/FEI	Business Operations
I World Pharmaceutical Co., Ltd.		688222857	manufacture(73442-0007)

Revised: 11/2019

I World Pharmaceutical Co., Ltd.