

WOMENS LAXATIVE- bisacodyl tablet
QUALITY CHOICE (Chain Drug Marketing Association)

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Bisacodyl USP 5 mg

Purpose

Stimulant laxative

Use

- for temporary relief of occasional constipation and irregularity
- this product generally produces a bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take with a glass of water

adults and children 12 years of age and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor

Other information

- each tablet contains: **calcium 15 mg**
- store between 20^o-25^oC (68^o-77^oF)

Inactive ingredients

acacia, calcium sulfate anhydrous, carnauba wax, colloidal silicon dioxide, corn starch, D&C red # 7 calcium lake, hypromellose phthalate, iron oxide, iron oxide black, iron oxide yellow, kaolin, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide, and triacetin

Questions or Comments?

Call **1-248-449-9300 Monday-Friday 9AM-5PM EST**

Principal Display Panel

Compare to the active ingredient in Correctol®*

Enteric Coated

Women's Laxative

Stimulant laxative

Bisacodyl, 5 mg

Gentle enough for sensitive stomachs

Tablets

*This product is not manufactured or distributed by MSD Consumer Care Inc., distributor of Correctol®.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BORKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATON.

Distributed by C.D.M.A., Inc. ©

43157 W. Nine Mile

Novi, MI 48376-0995

www.qualitychoice.com

Product Label

V60X4

PRINT SIDE GLUE



Directions

- Take with a glass of water

Drug Facts (continued)

■ a sudden change in bowel habits that lasts more than 2 weeks

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Questions or comments?
Call 1-248-449-9300 Monday-Friday 9AM-5PM EST

Inactive ingredients acacia, calcium sulfate anhydrous, carnauba wax, colloidal silicon dioxide, corn starch, D&C red #7 calcium lake, hypromellose pthalate, iron oxide, iron oxide black, iron oxide yellow, kaolin, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide, triacetin

Other information

- each tablet contains: calcium 15 mg
- store between 20°-25°C (68°-77°F)

Drug Facts (continued)

adults and children 12 years of age and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor

Drug Facts

Active ingredient (in each tablet) Bisacodyl USP 5 mg.....Stimulant laxative

Purpose

- this product generally produces a bowel movement in 6 to 12 hours

Warnings

- Do not use if you cannot swallow without chewing.
- Ask a doctor before use if you have stomach pain, nausea or vomiting.

Use

- for temporary relief of occasional constipation and irregularity

Drug Facts (continued)

NDC 63868-209-30



*Compare to the active ingredient in Correctol®

Enteric Coated
Women's Laxative
Stimulant Laxative

Enteric Coated Women's Laxative

Stimulant Laxative

Bisacodyl, 5 mg
Gentle Enough For Sensitive Stomachs



30 Tablets

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot No.:
Exp. Date:

*This product is not manufactured or distributed by MSD Consumer Care Inc., distributor of Correctol®.



Distributed by C.D.M.A., Inc.©
43157 W. Nine Mile
Novi, MI 48376-0995
www.qualitychoice.com
Questions: 248-449-9300



PLD-A14M
FC002388

Quality Choice Enteric Coated Women's Laxative

WOMENS LAXATIVE

bisacodyl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-209
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C RED NO. 7 (UNII: ECW0LZ41X8)	
HYPROMELLOSE PHTHALATE (24% PHTHALATE, 55 CST) (UNII: 87Y6436BKR)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
KAOLIN (UNII: 24H4NWX5CO)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE (UNII: FZ989GH94E)	
SHELLAC (UNII: 46N107B71O)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	TCL057
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-209-30	30 in 1 CARTON	08/31/2014	05/31/2024
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	08/31/2014	05/31/2024

Labeler - QUALITY CHOICE (Chain Drug Marketing Association) (011920774)

Revised: 11/2021

QUALITY CHOICE (Chain Drug Marketing Association)