

ACETAMINOPHEN- acetaminophen tablet
FAMILY DOLLAR

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Extra Strength
RAPID PAIN RELIEF
Acetaminophen USP, 500 mg
Pain reliever/Fever reducer

Active ingredient

(in each Gelcap)

Acetaminophen, USP 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- toothache
- the common cold
- premenstrual and menstrual cramps
- temporarily reduces fever

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
liver disease.

Ask a doctor or pharmacist before use if you are
taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,
ask a health professional before use.

Keep out of the reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed

adults and children 12 years and over

- take 2 gelcaps every 6 hours while symptoms last
- do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
- do not take more than 10 days unless directed by a doctor

children under 12 years

- ask a doctor

Other information

- store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature

- avoid high humidity
- see end panel for lot number and expiration date

Inactive ingredients


ammonium hydroxide, black iron oxide, black iron oxide irradiated, colloidal silicon dioxide, croscarmellose sodium, gelatin, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, hypromellose, iron oxide red, isopropyl alcohol, n-butyl alcohol, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, stearic acid, titanium dioxide, yellow iron oxide.

Questions or comments?

call **1-877-770-3183** Mon-Fri 9:00 AM to 4:30 PM EST.

Acetaminophen, USP 500 mg Rapid Release Gelcaps

SIDE 2



Finished Lead Edge

Drug Facts (continued)

Keep out of reach of children. If you are pregnant, trying to get pregnant, or breastfeeding, ask a health professional before use. Many out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Proper medical attention is critical for adults and may be fatal for children. Do not use for 10 days after surgery or anesthesia.

Warnings

Do not use if you have been advised by a doctor to avoid aspirin. Do not use if you have been advised by a doctor to avoid aspirin. Do not use if you have been advised by a doctor to avoid aspirin. Do not use if you have been advised by a doctor to avoid aspirin.

Directions

Take 2 capsules every 6 hours with meals and children 12 years and over. Do not take more than 6 capsules in 24 hours. Do not take more than 10 days unless directed by a doctor. Do not take more than 10 days unless directed by a doctor.

Other important information


Keep out of reach of children. Do not use if you have been advised by a doctor to avoid aspirin. Do not use if you have been advised by a doctor to avoid aspirin. Do not use if you have been advised by a doctor to avoid aspirin.

Contains 100 capsules


Questions or comments?

Call 1-877-378-3188 Mon-Fri 8:00 AM to 4:00 PM CST.

UVGS



SIDE 1



Finished Lead Edge

Family wellness
Extra Strength RAPID PAIN RELIEF
 Acetaminophen USP, 500 mg
 Pain Reliever/Fever Reducer
 100 Gelscaps

Drug Facts

Active Ingredient
 Acetaminophen USP, 500 mg (in each gelscap)

Pain relief/fever reducer

Warnings

Do not use if you have been advised by a doctor to avoid aspirin. Do not use if you have been advised by a doctor to avoid aspirin. Do not use if you have been advised by a doctor to avoid aspirin.

Directions

Take 2 capsules every 6 hours with meals and children 12 years and over. Do not take more than 6 capsules in 24 hours. Do not take more than 10 days unless directed by a doctor.

Other important information

Keep out of reach of children. Do not use if you have been advised by a doctor to avoid aspirin. Do not use if you have been advised by a doctor to avoid aspirin. Do not use if you have been advised by a doctor to avoid aspirin.

Contains 100 capsules

Questions or comments?

Call 1-877-378-3188 Mon-Fri 8:00 AM to 4:00 PM CST.

↑ .063" or

item	: LRU GRABOX 100C	view	: PRTVW
code #	:	date	: 08/7/10
size	: 1x3/4 x 1x3/4 x 3 1/2		
ref #	: 09180815		
material	: 016 085		

R-00
53200012
300000115
[Barcode]

Drug Facts (continued)

Warnings:
 • If you have ever had an allergic reaction to this product or any of its ingredients.
 • Ask a doctor or pharmacist before use if you have liver disease.
 • Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
 • Stop use and ask a doctor if:
 • pain gets worse or lasts more than 10 days
 • fever gets worse or lasts more than 3 days
 • new symptoms occur
 • redness or swelling is present
 • These could be signs of a serious condition.
 • If pregnant or breast-feeding, ask a health professional before use.
 • Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions:
 • Do not take more than directed.
 • Adults and children 12 years and over:
 • take 2 capsules every 6 hours while symptoms last.
 • Do not take more than 6 capsules in 24 hours, unless directed by a doctor.
 • Do not take more than 10 days unless directed by a doctor.
 • Children under 12 years:
 • ask a doctor.

Other information:
 • Store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature.

Drug Facts (continued)

Warnings:
 • Do not use in high humidity or in areas with high humidity for lot number and expiration date.
Inactive ingredients: aluminum hydroxide, black iron oxide, black iron oxide traditional, colloidal silicon dioxide, croscarmellose sodium, gelatin, hydroxypropyl cellulose, hydroxypropyl methylcellulose, hypromellose, iron oxide red, lactopropyl alcohol, *n*-butyl alcohol, polyethylene glycol, povidone, polyvinylpyrrolidone, propylene glycol, shellac, stearic acid, titanium dioxide, yellow iron oxide.

Questions or comments?
 • Call 1-877-776-3163 Mon-Fri 8:00 AM to 4:30 PM EST.

© All trademarks are property of their respective owners. This product is not affiliated with the manufacturers of Extra Strength Tylenol® Rapid Release Gels.

Distributed by:
 Millimed Brands, LLC
 500 Volvo Parkway
 Chesapeake, VA 23320

NOT 100% SATISFIED?
 Return package and unused product within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

Made in India

Extra Strength
RAPID PAIN RELIEF
 Acetaminophen USP, 500 mg
 Pain Reliever/Fever Reducer
FAMILY WELLNESS™

50 Gelcaps
Extra Strength RAPID PAIN RELIEF
 Acetaminophen USP, 500 mg Pain Reliever/Fever Reducer
FAMILY WELLNESS™
 CONTAINS 100 TYLENOL® EXTRA STRENGTH RAPID RELEASE GELS
 100% SATISFACTION GUARANTEED
 OTC
 Non-Prescription
 [Image of two capsules]

SKU 019884
 [Barcode]
 0 52251 52302 1
 Lot
 Exp.

Important: Read all product information before using.
 Drug facts contain the following information:
Warnings: Do not use if pregnant. SAFETY SEAL, UNDO CAP IS BROKEN OR MISSING.

Drug Facts

Active ingredient (in each gelcap)	Purpose
Acetaminophen, USP 500 mg...	Pain reliever/fever reducer

Uses:
 • temporarily relieves minor aches and pains due to:
 • headache
 • muscle aches
 • backache
 • minor pain of arthritis
 • the common cold
 • toothache
 • menstrual and menstrual cramps
 • temporarily reduces fever

Warnings:
Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take:
 • more than 4,000 mg of acetaminophen in 24 hours
 • with other drugs containing acetaminophen
 • 3 or more alcoholic drinks every day while using this product.
 Always alert acetaminophen may cause severe side reactions. Symptoms may include:
 • skin redness
 • hives
 • rash
 • If a skin reaction occurs, stop use and seek medical help right away.
 Do not use with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

31

item :	1BU GRABOX 100CN	view :	PRINT
code # :		date :	09/12
size :	1x3/4 x 1x3/4 x 3 1/2		
ref # :	FF180815		
material :	016 GRS		



R-00
300000115
532000012

Drug Facts (continued)

Warnings

- Do not take more than directed.
- Do not take more than 6 packages in 24 hours unless directed by a doctor.
- Do not take more than 10 days unless directed by a doctor.
- Ask a doctor if you have ever had an allergic reaction to this product or any of its ingredients.
- Ask a doctor before use if you have liver disease.
- Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
- Stop use and ask a doctor if you have pain worse or lasts more than 10 days or fever gets worse or lasts more than 3 days or new symptoms occur or redness or swelling is present.
- These could be signs of a serious condition.
- If pregnant or breast-feeding, ask a health professional before use.
- Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- adults and children 12 years and over: take 2 gelcaps every 6 hours while symptoms last. do not take more than 6 packages in 24 hours unless directed by a doctor.
- children under 12 years: ask a doctor.

Other information

Store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature.

Drug Facts (continued)

Warnings

- Do not take more than directed.
- Do not take more than 6 packages in 24 hours unless directed by a doctor.
- Do not take more than 10 days unless directed by a doctor.
- Ask a doctor if you have ever had an allergic reaction to this product or any of its ingredients.
- Ask a doctor before use if you have liver disease.
- Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
- Stop use and ask a doctor if you have pain worse or lasts more than 10 days or fever gets worse or lasts more than 3 days or new symptoms occur or redness or swelling is present.
- These could be signs of a serious condition.
- If pregnant or breast-feeding, ask a health professional before use.
- Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- adults and children 12 years and over: take 2 gelcaps every 6 hours while symptoms last. do not take more than 6 packages in 24 hours unless directed by a doctor.
- children under 12 years: ask a doctor.

Other information

Store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature.

Drug Facts (continued)

Warnings

- Do not take more than directed.
- Do not take more than 6 packages in 24 hours unless directed by a doctor.
- Do not take more than 10 days unless directed by a doctor.
- Ask a doctor if you have ever had an allergic reaction to this product or any of its ingredients.
- Ask a doctor before use if you have liver disease.
- Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
- Stop use and ask a doctor if you have pain worse or lasts more than 10 days or fever gets worse or lasts more than 3 days or new symptoms occur or redness or swelling is present.
- These could be signs of a serious condition.
- If pregnant or breast-feeding, ask a health professional before use.
- Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- adults and children 12 years and over: take 2 gelcaps every 6 hours while symptoms last. do not take more than 6 packages in 24 hours unless directed by a doctor.
- children under 12 years: ask a doctor.

Other information

Store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature.

Drug Facts

Active ingredient *Purpose*
(in each gelcap) *(for each gelcap)*
Acetaminophen, USP 500 mg... Pain reducer/fever reducer

Uses

- temporarily relieve minor aches and pains due to:
 - headache
 - muscular aches
 - inches
 - minor pain of arthritis
 - the common cold
 - flu
 - menstrual and menstrual cramps
 - temporarily reduce fever

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Other warnings: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blistering
- itching
- swelling
- difficulty breathing

If a skin reaction occurs, stop use and seek medical help right away.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

300000115
532000012
R-00

SKU 919884
0 52251 52302 1

Lot
Exp.

31"

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-167
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
AMMONIA (UNII: 5138Q19F1X)	
GELATIN (UNII: 2G86QN327L)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
BUTYL ALCOHOL (UNII: 8PJ61P6TS3)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	gray (Encapsulated with red opaque and blue gray opaque hard gelatin shells)	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	G1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-167-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	01/31/2020	
2	NDC:55319-167-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/31/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	01/31/2020	

Labeler - FAMILY DOLLAR (024472631)

Revised: 5/2020

FAMILY DOLLAR