

## **MILK OF MAGNESIA- magnesium hydroxide suspension**

### **Walgreens**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Drug Facts**

#### **Active Ingredients**

Magnesium hydroxide 1200 mg

#### **Purpose**

Saline laxative

#### **Uses**

relieves occasional constipation (irregularity). This product usually produces bowel movement in 1/2 to 6 hours

#### **Warnings**

##### **Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

##### **Ask a doctor or pharmacist before use if you are**

taking a prescription drug. This product may interact with certain prescription drugs.

##### **Stop use and ask a doctor if**

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you use a laxative for more than 1 week

##### **If pregnant or breast-feeding**

ask a health professional before use.

##### **Keep out of reach off children**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### **Directions**

- do not exceed the maximum recommended daily dose in a 24-hour period
- shake well before using
- mL = milliliter
- measure only with dosing cup provided. do not use any other dosing device.

- keep dosing cup with product
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor. Drink a full glass (8 oz) of liquid with each dose.

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	do not use

### Other information

- each 15 mL contains: magnesium 500 mg
- store between 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). Avoid freezing. Close cap tightly after use.

### Inactive Ingredients

purified water, sodium hypochlorite

### Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

### Principal Display Panel

Compare to Phillips'® Milk of Magnesia active ingredient<sup>††</sup>

### Milk of Magnesia

MAGNESIUM HYDROXIDE 1200 mg / SALINE LAXATIVE

SUGAR FREE

STIMULANT FREE

CRAMP FREE

- Comfortable, dependable overnight relief

ORIGINAL FLAVOR

This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips'® Milk of Magnesia.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

**DISTRIBUTED BY: WALGREEN CO.**

**200 WILMOT RD., DEERFIELD, IL 60015**

walgreens.com

### Package label

**VALUE SIZE**

*Walgreens*

Compare to Phillips® Milk of Magnesia active ingredient<sup>11</sup>

NDC 0363-0847-26

# Milk of Magnesia

MAGNESIUM HYDROXIDE 1200 mg / SALINE LAXATIVE

**SUGAR FREE**    **STIMULANT FREE**

**CRAMP FREE**

• Comfortable, dependable overnight relief

ORIGINAL FLAVOR

26 FL OZ (1 PT 10 OZ) 769 mL

PLD-A454A  
LB005698

**Drug Facts**

Active ingredient (in each 15 mL)	Purpose
Magnesium hydroxide 1200 mg..... Saline laxative	

**Use** Relieves occasional constipation (irregularity). This product usually produces bowel movement in 1/2 to 6 hours.

**Warnings**

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug. This product may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not exceed the maximum recommended daily dose in a 24-hour period
- shake well before using
- mL = milliliter
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product

**Drug Facts (continued)**

■ dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor. Drink a full glass (8 oz) of liquid with each dose.

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	do not use

**Other information**

- each 15 mL contains: magnesium 500 mg
- store between 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). Avoid freezing. Close cap tightly after use.

**Inactive ingredients**  
purified water, sodium hypochlorite


**Questions or comments?** Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING**

<sup>11</sup>This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips® Milk of Magnesia.

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2018 Walgreen Co.

ITEM 290106 W10023-0418-F



3 11917 20021 7

PLD-A454A LB005698 ORG0318-F

**WALGREENS Milk of Magnesia**

**MILK OF MAGNESIA**

magnesium hydroxide suspension

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-0847
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

**Inactive Ingredients**

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0847-12	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2018	
2	NDC:0363-0847-26	769 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2018	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part334	03/31/2018	

**Labeler** - Walgreens (008965063)

Revised: 10/2019

Walgreens