

**ACETAMINOPHEN 325MG- acetaminophen tablet**  
**Pharmaceutica North America, Inc.**

-----

**Active Ingredient(in each tablet)**

Acetaminophen 325mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children.

Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if the user has** liver disease

**Ask a doctor or pharmacist before use if the user is**taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years

- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of the reach of children.**

**Overdose warning:**

In the case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

do not take more than directed (see overdose warning)

---

adults and children 12 years and over	<ul style="list-style-type: none"> <li>• take a 2 tablets, every 4 to 6 hours while symptoms last</li> <li>• do not take more than 10 tablets in 24 hours, unless direct by a doctor</li> <li>• do not use more than 10 days unless directed by a doctor</li> </ul>
children 6 to under 12 years	<ul style="list-style-type: none"> <li>• take a one tablet, every 4 to 6 hours while symptoms last</li> <li>• do not take more than 5 tablets in 24 hours</li> <li>• do not use more than 5 days unless directed by a doctor</li> </ul>
children under 6 years	ask a doctor

---

**Other information**

- store between 20-25°C (68-77°F)

**Inactive Ingredients**

povidone, pregelatinized corn starch, sodium starch glycolate, stearic acid

**Questions or comments? (888)788-6472**

**Product label**



NDC 45861-085-01

# Acetaminophen

## Regular Strength

### 325mg Tablet

Pain Reliever / Fever Reducer

1000 Tablets

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**Drug Facts**  
**Active Ingredient (in each tablet)**  
 Acetaminophen 325mg

**Purpose**  
 Pain reliever / fever reducer

**Uses**  
 temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ backache ■ minor pain of arthritis ■ the common cold ■ toothache ■ premenstrual and menstrual cramps ■ temporarily reduces fever

**Warnings**  
 Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if adult takes more than 4,000 mg of acetaminophen in 24 hours child takes more than 5 doses in 24 hours, which is the maximum daily amount taken with other drugs containing acetaminophen adult has 3 or more alcoholic drinks every day while using this product  
 Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin redness ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription).  
 ■ If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ If you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if the user has liver disease**

**Stop use and ask doctor if** ■ pain gets worse or lasts more than 10 days in adults ■ pain gets worse or lasts more than 5 days in children under 12 years ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present  
 These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**  
**Overdose warning:** In the case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

■ do not take more than directed (see overdose warning)  
 adults and 12 years and over take 2 tablets, every 4 to 6 hours while symptoms last do not use for more than 10 days unless directed by a doctor  
 children 6 years and over take 1 tablet every 4 to 6 hours while symptoms last do not take more than 5 tablets in 24 hours  
 children under 12 years do not use for more than 5 days unless directed by a doctor  
 children under 6 years ask a doctor

**Other information**  
 store between 20°-25°C (68°-77°F)

**Questions or comments? (888)788-6472**

Manufactured For:  
 Pharmacia North America, Inc.  
 Glendale, CA 91205



## ACETAMINOPHEN 325MG

acetaminophen tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:45861-085
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

### Product Characteristics

Color	white (Off White)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:45861-085-	1000 in 1 BOTTLE; Type 0: Not a Combination	02/01/2025	

01	Product	02/01/2025	
<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/01/2025	

**Labeler** - Pharmaceutica North America, Inc. (962739699)

Revised: 3/2025

Pharmaceutica North America, Inc.