QUICK-SAN- sanitizing teat dip liquid IBA

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Quick-San

Active Ingredients

ACTIVE INGREDIENTS:

Hydrogen Peroxide	0.60%
Lactic Acid	0.50%

INERT INGREDIENTS:

Glycerin	.2.00%
Other InertsL	.96.90%
(Includes Buffering Agents, Stabilizing Agents, Wetting	g Agents
and Colorants)	
TOTAL	L00.00%

GENERAL RECOMMENDATION

GENERAL RECOMMENDATION

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.

2. When solution in the teat dip cup becomes visibly dirty, discard remaining contents and refill cup with fresh teat dip.

3. Do not use this product for cleaning or sanitizing equipment.

4. Wash teat dip cups after each milking.

CAUTION

KEEP OUT OF REACH OF CHILDREN. AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY. PROTECT EYES AND MUCOUS MEMBRANES FROM CONTACT WITH THIS PRODUCT.

FIRST AID PROCEDURES FIRST AID PROCEDURES **EYE CONTACT**: AVOID CONTACT WITH EYES. IF CONTACT OCCURS, FLUSH EYE WITH LARGE QUANTITIES OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUE. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE. IF IRRIGATION DEVELOPS, CONSULT A PHYSICIAN.

SKIN CONTACT: WASH ANY SENSITIVE AREAS.

INGESTION: DRINK TWO GLASSES OF WATER AND CONSULT A PHYSICIAN.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION

FOR MEDICAL EMERGENCIES CALL 1-800-424-9300

RECOMMENDED PROCEDURES FOR TEAT SANITATION

RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

1. Remove visible soils from the teats.

2. Observe foremilk by stripping two or three streams of milk into a strip cup.

3. Dip, foam or spray the cow's teats full length with **QUICK-SAN** Teat Dip.

4. Wait approximately 30 seconds.

5. Remove all teat dip by thoroughly drying the teats, teat orifices and udder (if foaming or spraying) with a clean, single, service paper towerl(s) to avoid contamination of milk.

6. Attach milking unit.

B. POST-MILKING (POST-DIPPING)

1. Immediately after milking, dip or spray all teats full length with **QUICK-SAN** Teat Dip. **Ensure good coverage of all sides of the teats and teat orifices.**

WARNING

WARNING

DO NOT MIX QUICK-SAN TEAT DIP WITH ANY OTHER PRODUCT.


DO NOT DILUTE THIS PRODUCT.

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.

ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.

PROTECT THIS PRODUCT FROM FREEZING.

Not For Human Use



QUICK-SAN

sanitizing teat dip liquid					
Product Information					
Product Type	OTC ANIMAL DRUG	Item Code (Source)		NDC:29983-1054	
Route of Administration	TOPICAL				
Active Ingredient/Active Moiety					
Ingredient Name			Basis of Strength		Strength
HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)			HYDROGEN PEROXIDE		.60 kg in 100 kg
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)		LACTIC ACID		.50 kg in 100 kg	

#	ltem Code	Package Description	Marketing	Start Date	Marketing	End Date
1	NDC:29983-1054-4	19.14 kg in 1 PAIL				
2	NDC:29983-1054-6	57.42 kg in 1 DRUM				
3	NDC:29983-1054-8	210.56 kg in 1 DRUM				
4	NDC:29983-1054-9	1052.78 kg in 1 CONTAINER				
Marketing Information						
	Marketing Category	Application Number or Mo Citation	onograph	Marketing S Date	tart Mark	eting End Date
	approved drug ner			09/27/2010		

Labeler - IBA (019494160)

Registrant - IBA (019494160)

Establishment					
Name	Address	ID/FEI	Business Operations		
Morgan Gallacher Inc. DBA Custom Chemical Formulators Inc.		028311595	manufacture, api manufacture, pack		

Revised: 12/2023

IBA