

**DG HEALTH IBUPROFEN- ibuprofen tablet, film coated
Dolgenercorp Inc**

Dolgenercorp, LLC Ibuprofen Drug Facts

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

Adults and children 12 years and older:

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor

Children under 12 years: ask a doctor

Other information

- read all warnings and directions before use
- store between 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- see end panel for lot number and expiration date

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide

Questions or comments?

1-888-309-9030

Principal Display Panel

DG™ |health

SEE REVISED WARNING

Compare to the active ingredient of Advil® Ibuprofen Caplets

Ibuprofen Tablets, 200 mg

Pain Reliever/Fever Reducer(NSAID)

50 Caplets**

**Capsule-Shaped Tablets

200 mg

Actual Caplet Size



Important: Read all product information before use. Keep this box for important information.

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING



Actual Size

DG health
Ibuprofen
Tablets, 200 mg
Pain Reliever/
Fever Reducer
(NSAID)

50 Caplets**
**Capsule-Shaped Tablets

Drug Facts

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)..... Pain reliever/fever reducer/anti-inflammatory drug

Purposes

■ temporarily relieve minor aches and pains due to:
■ headache ■ muscle aches ■ minor pain of arthritis ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever

Warnings

■ Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin rash ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.

■ Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed ■ have had a stroke ■ take aspirin, aspirin/acetaminophen, or other heart attack and stroke warning NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as:

Drug Facts (continued)

as heartburn ■ you have high blood pressure, heart disease, heart failure, kidney disease, asthma, or had a stroke ■ you are taking a diuretic. Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug.

When using this product

■ Take with food or milk if stomach upset occurs. Stop use and ask a doctor if ■ you experience any of the following signs or symptoms: ■ heartburn ■ vomiting ■ dark stools ■ black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain get worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the paranasal area ■ any new symptoms appear.

If pregnant or breast-feeding, ask a health professional before use. It is especially important to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-272-1222).

Directions ■ do not take more than directed ■ the smallest effective dose should be used. Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms persist. ■ If pain or fever does not respond to 1 caplet, 2 caplets may be used. ■ do not exceed 2 caplets in 24 hours, unless directed by a doctor. Children under 12 years: ask a doctor.

100% Satisfaction Guarantee!

**This product is not manufactured or distributed by GSK CH, distributor of Advil® Ibuprofen Caplets. DISTRIBUTED BY OLEAST MAN CO. 10 MISSOURI RD. GODFREY, TN 37072 (888) 309-9030

Questions or comments?

■ **mac drug ingredients** color bit silicon dioxide, corn starch, croscarmellose sodium, hypromellose, polyacrylate 80, polyethylene glycol, polyacrylate 80, stearic acid, titanium dioxide
■ **Other information** ■ read all warnings and directions before use ■ store below 20-25°C (68-77°F) ■ avoid high humidity and excessive heat above 40°C (104°F) ■ see end panel for lot number and expiration date
■ **Questions or comments?** 1-888-309-9030

SEE REVISED WARNING
Compare to the active ingredient of Advil® Ibuprofen Caplets**

DG health

Ibuprofen

Tablets, 200 mg
Pain Reliever/Fever Reducer
(NSAID)

50 Caplets**



200 mg

**Capsule-Shaped Tablets
Actual Caplet Size

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-647
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	BROWN	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	I2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-647-78	1 in 1 CARTON	06/17/2021	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55910-647-71	1 in 1 CARTON	06/16/2021	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
---------------------------	---	-----------------------------	---------------------------

ANDA	ANDA072096	06/16/2021	
------	------------	------------	--

Labeler - Dolgencorp Inc (068331990)

Revised: 1/2024

Dolgencorp Inc