LIFT EYECREAM- glycerin cream ROA Pacific Corporation

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Drug Facts

glycerin

Water, Propylene Glycol, Sodium Hyaluronate, Butylene Glycol, Glyceryl Stearate, Castanea Crenata(Chestnut) Shell Extract, Centella Asiatica Extract, Hydrolyzed Collagen, Xanthan Gum, Carbomer, Triethanolamine, PEG-60 Hydrogenated Castor Oil, Dipalmitoyl Hydroxyproline, Chlorphenesin, Glycyrrhiza Uralensis (Licorice) Extract, Linum Usitatissimum(Linseed) Seed Extract, Hibiscus Esculentus Fruit Extract, Aloe Barbadensis Leaf Juice, Arctium Lappa Root Extract, Hibiscus Mutabilis Flower Extract, Corchorus Olitorius Leaf Extract, Camellia Sinensis Leaf Extract, Portulaca Oleracea Extract, Perfume, Red Oxide of Iron, Adenosine, Magnesium Aluminum Silicate, Panthenol, Sodium DNA, Glycine HCl, Proline, Sodium Ascorbyl Phosphate, Acetyl Methionine, Theanine, Acetyl Hexapeptide-8, Disodium EDTA, Allantoin, Olea Europaea (Olive) Fruit Oil, Phytosterols, Vegetable Squalene, Butyrospermum Parkii (Shea) Butter, Ceramide3, Propanediol

-The instant botox effect lasts average 5-8hours just like a magic of two minutes.

-Great effects on skin moisturizing and pore tightening

-Overall lifting effect when it is applied to the whole face

-Non-sticky and smooth texture

keep out of reach of the children

• After cleansing your face, take a small amount of it and then massage the eye rims circlewise from inside to outside. After massaging repetitively 2-3 times, put on the basic skincare products.

[Step1] After cleansing your face, wipe any fluids and then use it before the basic skincare.

[Step2] Take a suitable amount on the back of your hand.

[Step3] Massage the eye rims circlewise from inside to outside.

[Step4] After the cream is fully absorbed(20-30 seconds later), apply the cream left on the back of your hand 2-3 times repetitively.


■ if following abnormal symptoms occurs after use , stop use and consult with a skin specialist

red specks, swelling, itching

■ don't use on the part where there is injury, eczema, or dermatitis

Keep out of reach of children

■ if swallowed, get medical help or contact a person control center immediately for external use only



LIFT EYECRE	AM						
glycerin cream							
Product Inform	ation						
Product Type		HUMAN OTC DRUG Item Code (Source)				NDC:73490-0001	
Route of Administ	tration	TOPICAL					
Active Ingredient/Active Moiety							
Ingredient Name Basis of S					gth	Strength	
GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX)				GLYCERIN 5 g i		5 g in 100 mL	
Inactive Ingred	ionto						
Inactive Ingredients							
Ingredient Name WATER (UNII: 059QF0K00R)					•	Strength	
BUTYLENE GLYCOL		KORA)					
Packaging							
# Item Code	Pa	Package Description		Marketing Start Date		Marketing End Date	
- NDC:73490- 1		TTLE; Type 0: Not a Combi					

• 0001-1	Product	11/20/2019				
Marketing Information						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
unapproved drug		11/16/2019				
other						

Labeler - ROA Pacific Corporation (631167348)

Registrant - ROA Pacific Corporation (631167348)

Establishment							
Name	Address	ID/FEI	Business Operations				
ROA Pacific Corporation		631167348	manufacture(73490-0001) , label(73490-0001)				

Revised: 10/2022

ROA Pacific Corporation