

QUICK-SAN 10X CONCENTRATE- non-iodine sanitizing teat dip liquid IBA

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).

Quick-San 10X Concentrate

ACTIVE INGREDIENT

Hydrogen Peroxide.....6.00%

(Provides A Full .6% Hydrogen Peroxide When Mixed At The Recommended Dilution Ratio)

Lactic Acid.....5.00%

(Provides A Full .5% Lactic Acid When Mixed At The Recommended Dilution Ratio)

INERT INGREDIENTS

Glycerin.....20.00%

(Provides A Total 2% Glycerin At The Recommended Dilution Ratio)

Other Inerts.....69.00%

(Includes Buffering Agents, Stabilizing Agents, and Surfactants)

TOTAL.....100.00%

GENERAL RECOMMENDATIONS

A.) A quality water supply is required for mixing.

- 1.) Hardness <2 grains or 35 ppm
- 2.) Iron must be 0.0 ppm
- 3.) pH 6.5-7.0

B.) CONCENTRATE should be mixed into ready-to-use product within 30 days of delivery,

C.) Keep CONCENTRATE out of direct sunlight and store in a cool place

D.) QUICK-SAN 10X CONCENTRATE contains a high level of Hydrogen Peroxide and should be handled with care. This concentration can be detrimental to the skin, eyes and soft tissues.

FIRST AID PROCEDURES

IF IN EYES: RINSE CAUTIOUSLY WITH WATER FOR SEVERAL MINUTES. REMOVE CONTACT LENSES, IF PRESENT, AND EASY TO DO. CONTINUE RINSING WITH WATER FOR 15 MINUTES. GET IMMEDIATE MEDICAL ADVICE/ATTENTION.

IF SWALLOWED: RINSE MOUTH WITH WATER. DO NOT INDUCE VOMITING. GET IMMEDIATE MEDICAL ADVICE/ATTENTION.

IF ON SKIN: IMMEDIATELY REMOVE ALL CONTAMINATED CLOTHING AND WASH BEFORE RE-USE. WASH SKIN WITH PLENTY OF WATER FOR 15 MINUTES. IF SKIN IRRITATION OCCURS, GET MEDICAL ADVICE/ATTENTION

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300

MIXING GUIDELINES FOR QUICK- SAN 10X TEAT DIP CONCENTRATE

1. Clean mixing equipment, clean transfer lines, clean and sanitize packing containers.
2. Use only equipment that is compatible with Hydrogen Peroxide and Acids.
3. Start with QUICK- SAN 10X CONCENTRATE.
4. Water volume to QUICK- SAN 10X CONCENTRATE ratio at least 5:1.
5. With agitation, slowly add the QUICK- SAN 10X CONCENTRATE to the water.
6. Use additional water to rinse QUICK- SAN 10X CONCENTRATE residue from QUICK- SAN 10X CONCENTRATE container and add to mixture.
7. Continue mixing to a uniform blend.
8. Continue mixing while adding the balance of water or skin conditioners to the desired concentration.
9. Test for Hydrogen Peroxide strength pH.
10. Assign a lot number , record test results and retain a sample of finished product.
11. Fill properly labeled containers suitable for Hydrogen Peroxide based products.

CAUTION: KEEP OUT OF REACH OF CHILDREN, STORE LOCKED UP.
AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

DILUTE 1 PART
QUICK- SAN 10X TEAT DIP
CONCENTRATE
Per 9 Parts Potable,
Tepid Water.

AN AID IN THE
PREVENTION OF NEW
INTRAMAMMARY
INFECTIONS

WARNING

- QUICK-SAN 10X NPE FREE TEAT DIP CONCENTRATE MUST BE DILUTED BEFORE USE. DO NOT MIX WITH ANYTHING BUT POTABLE WATER.
- WEAR SAFETY EQUIPMENT. A MINIMUM OF SAFETY GOGGLES AND RUBBER GLOVES.
- VENTED BUNGS ARE REQUIRED ON THE CONTAINER OF READY-TO-USE PRODUCT.
- SECURE BUNGS/CAPS TO ENSURE THE PRODUCT INTEGRITY.
- USE DEDICATED EQUIPMENT FOR BLENDING QUICK-SAN 10X CONCENTRATE. DO NOT USE EQUIPMENT THAT HAS CONTACT WITH OTHER CHEMICALS.
- DO NOT LET THE QUICK-SAN 10X CONCENTRATE OR READY-TO-USE PRODUCT COME INTO CONTACT WITH BLACK STEEL AND IRON, SOFT METALS, ALLOYS OF

SOFT METALS, WOODOR PAPER.

- DISPOSE OF CONTENTS/CONTAINER IN ACCORDANCE WITH LOCAL, STATE AND FEDERAL REGULATIONS.
- PROTECT THIS PRODUCT FROM FREEZING

Quick-San 10X Concentrate

NDC 29983-1070-9

THE
SHIELD
OF

IBA

QUALITY
AND
VALUE

QUICK-SAN 10X CONCENTRATE

A DUAL GERMICIDE, NPE FREE, NON-IODINE TEAT DIP

ACTIVE INGREDIENTS:

Hydrogen Peroxide..... 6.00%
(Provides A Full .6% Hydrogen Peroxide When Mixed
At The Recommended Dilution Ratio)

Lactic Acid 5.00%
(Provides a Full .5% Lactic Acid When Mixed
At The Recommended Dilution Ratio)

INERT INGREDIENTS:

Glycerin.....20.00%
(Provides a total of 2% Glycerin At The Recommended Dilution Ratio)

Other Inerts.....69.00%
(Includes Buffering Agents, Stabilizing Agents and Surfactants)

TOTAL:..... 100.00%

GENERAL RECOMMENDATIONS

A.) A quality water supply is required for mixing.

1.) Hardness < 2 grains or 35 ppm.

2.) Iron must be 0.0 ppm.

3.) pH 6.5 - 7.0.

B.) **CONCENTRATE** should be mixed into ready-to-use product within 30 days of delivery.

C.) Keep **CONCENTRATE** out of direct sunlight and store in a cool place.

D.) **QUICK-SAN 10X CONCENTRATE** contains a high level of Hydrogen Peroxide and should be handled with care. This concentration can be detrimental to the skin, eyes and soft tissues.

CAUTION: KEEP OUT OF REACH OF CHILDREN. STORE LOCKED UP. AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

FIRST AID PROCEDURES

IF IN EYES: RINSE CAUTIOUSLY WITH WATER FOR SEVERAL MINUTES. REMOVE CONTACT LENSES, IF PRESENT, AND EASY TO DO. CONTINUE RINSING WITH WATER FOR 15 MINUTES. GET IMMEDIATE MEDICAL ADVICE/ATTENTION.

IF SWALLOWED: RINSE MOUTH WITH WATER. DO NOT INDUCE VOMITING. GET IMMEDIATE MEDICAL ADVICE/ATTENTION.

IF ON SKIN: IMMEDIATELY REMOVE ALL CONTAMINATED CLOTHING AND WASH BEFORE RE-USE. WASH SKIN WITH PLENTY OF WATER FOR 15 MINUTES. IF SKIN IRRITATION OCCURS, GET MEDICAL ADVICE/ATTENTION.

**ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300**

**MIXING GUIDELINES FOR
QUICK-SAN 10X TEAT DIP CONCENTRATE**

1. Clean mixing equipment, clean transfer lines, clean and sanitize packaging containers.

2. Use only equipment that is compatible with Hydrogen Peroxide and Acids.

3. Start with **QUICK-SAN 10X CONCENTRATE**.

4. Water volume to **QUICK-SAN 10X CONCENTRATE** ratio at least 5:1.

5. With agitation, slowly add the **QUICK-SAN 10X CONCENTRATE** to the water.

6. Use additional water to rinse **QUICK-SAN 10X CONCENTRATE** residue from **QUICK-SAN 10X CONCENTRATE** container and add to mixture.

7. Continue mixing to a uniform blend.

8. Continue mixing while adding the balance of water or skin conditioners to the desired concentration.

9. Test for Hydrogen Peroxide strength and pH.

10. Assign a lot number, record test results and retain a sample of finished product.

11. Fill properly labeled containers suitable for Hydrogen Peroxide based products.

RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

1. Remove visible soils from the teats.

2. Observe foremilk by stripping two or three streams of milk into a strip cup.

3. Dip, foam or spray the cow's teats full length with QUICK-SAN Teat Dip.

4. Wait approximately 30 seconds.

5. Remove all teat dip by thoroughly drying the teats, teat orifices and udder (if foaming or spraying) with a clean, single service paper towel(s) to avoid contamination of milk.

6. Attach milking unit.

B. POST-MILKING (POST-DIPPING)

1. Immediately after milking, dip or spray all teats full length with QUICK-SAN Teat Dip. Ensure good coverage of all sides of the teats and teat orifices.

WARNING

- QUICK-SAN 10X NPE FREE TEAT DIP CONCENTRATE **MUST BE DILUTED BEFORE USE. DO NOT MIX WITH ANYTHING BUT POTABLE WATER.**
- WEAR SAFETY EQUIPMENT. A MINIMUM OF SAFETY GOGGLES AND RUBBER GLOVES.
- VENTED BUNGS ARE REQUIRED ON THE CONTAINER OF READY-TO-USE PRODUCT.
- SECURE BUNGS/CAPS TO ENSURE THE PRODUCT INTEGRITY.
- USE DEDICATED EQUIPMENT FOR BLENDING QUICK-SAN 10X CONCENTRATE. DO NOT USE EQUIPMENT THAT HAS CONTACT WITH OTHER CHEMICALS.
- DO NOT LET THE QUICK-SAN 10X CONCENTRATE OR READY-TO-USE PRODUCT COME IN CONTACT WITH BLACK STEEL AND IRON, SOFT METALS, ALLOYS OF SOFT METALS, WOOD OR PAPER.
- DISPOSE OF CONTENTS/CONTAINER IN ACCORDANCE WITH LOCAL, STATE AND FEDERAL REGULATIONS.
- PROTECT THIS PRODUCT FROM FREEZING.

Dilute 1 Part
**QUICK-SAN 10X TEAT DIP
CONCENTRATE**
Per 9 Parts Potable,
Tepid Water.

**AN AID IN THE
PREVENTION OF NEW
INTRAMAMMARY
INFECTIONS**

Not For Human Use

A PART OF THE COMPLETE IBA SANITATION PROGRAM

EXP. DATE: LOT NO.: NET CONTENTS: 250 GALLONS (946.353 LITERS) IBA STOCK # 261000 124W

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 103 GILMORE DRIVE, SUTTON, MA 01590

QUICK-SAN 10X CONCENTRATE

non-iodine sanitizing teat dip liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-1070
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE -	HYDROGEN	6 kg

UNII:BBX060AN9V)	PEROXIDE	in 100 kg
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)	LACTIC ACID	5 kg in 100 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-1070-7	114.26 kg in 1 DRUM		
2	NDC:29983-1070-9	1038.73 kg in 1 CONTAINER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		11/21/2017	

Labeler - IBA (019494160)

Registrant - IBA (019494160)

Establishment

Name	Address	ID/FEI	Business Operations
Morgan Gallacher Inc DBA Custom Chemical Formulator Inc		028311595	manufacture, api manufacture, pack