

ACETAMINOPHEN 250MG ASPIRIN 250MG CAFFEINE 65MG- acetaminophen 250mg aspirin 250mg caffeine 65mg tablet WALGREENS CO.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

**Walgreens Headache Relief Geltabs
Acetaminophen 250mg, Aspirin 250mg, Caffeine 65mg**

Acetaminophen 250mg

Aspirin 250mg

Caffeine 65mg

ammonium hydroxide, black iron oxide, brilliant blue FCF, corn starch, D&C yellow No. 10 lake, ethyl cellulose, FD&C blue No. 1 lake, FD&C red No. 40 lake, gelatin, glycerin, hydroxypropyl methyl cellulose, lactose monohydrate, light liquid paraffin, microcrystalline cellulose, povidone, pregelatinised starch, propylene glycol, shellac glaze, sorbitol sorbitan, stearic acid, talc, titanium dioxide, triacetin.

- Do not use more than directed
- Drink a full glass of water with each dose
- Adults and children 12 years and over: take 2 geltabs every 6 hours; not more than 8 geltabs in 24 hours
- Children under 12 years: ask a doctor

Temporarily relieves minor aches and pains due to:

Headache, a cold, arthritis, muscular aches, toothache, premenstrual and menstrual cramps

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing), shock.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 8 geltabs in 24 hours, which is the maximum daily amount, with other drugs contains acetaminophen, 3 or more alcoholic drinks every day while using this product.

Stomach bleeding warning: This contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take

other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using product, take more or for a longer time than direct.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/ fever, with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease, stomach bleeding warning applies to you, you have a history of stomach problems, such as heartburn, you have high blood pressure, heart disease, liver cirrhosis, or kidney disease, you are taking a diuretic, you have asthma.

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout, or arthritis, any other drug, or are under a doctor's care for any serious condition.

Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away. You experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, ringing in the ears or loss of hearing occurs, painful area is red or swollen, pain gets worse or lasts for more than 10 days, fever gets worse or lasts for more than 3 days, any new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Keep out of reach of children.

OVERDOSE WARNING: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away.

Pain reliever

Pain reliever

Pain reliever aid

- Store at room temperature between 20-25 °C (68-77 °F)
- Avoid excessive heat, cold, and humidity
- Close cap tightly after use
- Read all product information before using. Keep box for important information.

LEBG909M

Walgreens

Compare to Excedrin® Extra Strength active ingredients!

Headache Relief

NDC 0363-1320-23

ACETAMINOPHEN / ASPIRIN (NSAID) / CAFFEINE
PAIN RELIEVER / PAIN RELIEVER AID

EXTRA STRENGTH GELTABS

20 GELTABS



ACTUAL SIZE

CT132023 REV 122018
Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey
*This product is not manufactured or distributed by Novartis Consumer Health, Inc., owner of the registered trademark Excedrin® Extra Strength.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2018 Walgreen Co.
ITEM 129562 W10058-0519-L



3 11917 20713 1

Lot No.:
Exp. Date:

Drug Facts (continued)

Other information ■ store at room temperature between 20-25°C (68 to 77°F) ■ avoid excessive heat, cold and humidity ■ close cap tightly after use ■ read all product information before using. Keep box for important information.

Inactive ingredients Ammonium Hydroxide, Black Iron Oxide, Brilliant Blue FCF, Carn Starch, D&C Yellow No.10 Lake, Ethyl Cellulose, FD&C Blue No.1 Lake, FD&C Red No.40 Lake, Gelatin, Glycerin, Hydroxypropyl methyl cellulose, Lactose monohydrate, Light Liquid Paraffin, Microcrystalline Cellulose, Povidone, Prejelatinised Starch, Propylene Glycol, Shellac Glaze, Sorbitol Sorbitan, Stearic Acid, Talc, Titanium dioxide, Triacetin

Questions or comments? 1-888-677-8333



Drug Facts (continued)

■ any other drug, or are under a doctor's care for any serious condition
■ See your doctor if:
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ ringing in the ears or loss of hearing occurs
■ painless area is red or swollen
■ pain gets worse or lasts for more than 10 days
■ fever gets worse or lasts for more than 3 days
■ any new symptoms occur
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless expressly directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
OVERDOSE WARNING: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
■ do not use more than directed
■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 2 gel tabs every 6 hours; not more than 8 gel tabs in 24 hours
■ children under 12 years: ask a doctor

ORG0218-F

Drug Facts (continued)

■ 3 or more alcoholic drinks every day while using this product
Special bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ take 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.
Do not use ■ If you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver disease, or kidney disease ■ you are taking a diuretic ■ you have asthma
Ask a doctor or pharmacist before use if you are taking ■ a prescription drug for diabetes, gout, or arthritis

Drug Facts

Important information: THIS PRODUCT IS NOT FOR USE IN SEALS. IMPROPERLY SEALED BOTTLES MAY BE DAMAGED. "SEALING FOR YOUR PROTECTION" IS MARKED ON THE BOTTLE.

Active ingredients (in each gel tab)
Acetaminophen 320 mg...Pain reliever
Aspirin 320 mg (NSAID).....Pain reliever
Caffeine 65 mg.....Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses
Temporarily relieves minor aches and pains due to:
■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache ■ premenstrual & menstrual cramps

Warnings
Do not use: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Always alert: Acetaminophen may cause severe skin reactions.
Symptoms may include: ■ skin redness ■ blisters ■ rash ■ if a skin reaction occurs, stop use and seek medical help right away.
Always alert: Aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Use warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 8 gel tabs in 24 hours, which is the maximum daily amount, or with other drugs containing acetaminophen.

1 2 3 4 5
6 7 8 9
7690007
573728

Walgreens

Compare to Excedrin® Extra Strength active ingredients!

NDC 0385-1320-80

Headache Relief

ACETAMINOPHEN / ASPIRIN (NSAID) / CAFFEINE PAIN RELIEVER / PAIN RELIEVER AID

EXTRA STRENGTH GELTABS

80 GELTABS

ACTUAL SIZE



Drug Facts (continued)

Do not use if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease

- if you have a history of stomach problems, such as heartburn
- if you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- if you are taking a diuretic
- if you have asthma

Ask a doctor or pharmacist before use if you are taking

- a prescription drug for diabetes, pain, or arthritis
- any other drug, or are under a doctor's care for any serious condition

Show me and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - black stool
 - vomit blood
 - have bloody or black stools
- you have stomach pain that does not get better
- itching in the ears or loss of hearing occurs
- your skin is red or swollen
- your lips get worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition

If present or worsened, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

OVERDOSE WARNING: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- do not take more than 12 tablets in 24 hours
- do not take more than 8 tablets in 24 hours
- ask a doctor under 12 years; ask a doctor

Lot No. :
Exp. Date:

3 11917 20714 8

ITEM 128598 W10058-0519-L

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD, DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
Walgreens.com ©2018 Walgreen Co.
Walgreens Pharmacist Recommended
TradeMark Excedrin® Extra Strength.
Health, Inc., owner of the registered
distributor by Novartis Consumer
if this product is not manufactured or

Drug Facts

Active Ingredients (in each tablet)

- Acetaminophen 250 mg..... Pain reliever
- Aspirin 250 mg (NSAID)..... Pain reliever
- Caffeine 65 mg..... Pain reliever aid

Contains: nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - cold
 - arthritis
 - muscular aches
 - backache
 - premenstrual & menstrual cramps

Warnings

Heart's symptoms: Children and teenagers who have or are recovering from chicken pox or the flu symptoms should not use this product.

When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Stomach aches: Acetaminophen may cause severe skin reactions. If symptoms may include: skin rash, itching, hives, or skin peeling.

Aspirin: Aspirin may cause a severe allergic reaction, which may include: wheezing or trouble breathing, which may be fatal. If you have had asthma, ask your doctor before using this product because it may cause severe stomach bleeding. The chances are higher if you are age 60 or older.

Other warnings: This product contains acetaminophen. Severe liver damage may occur if you take more than 8 tablets in 24 hours, which is the maximum daily amount.

Do not use if you are taking

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you are age 60 or older.

- if you have had stomach ulcers or bleeding problems
- if you take a blood thinner (anticoagulant) or steroid drug
- if you take other drugs containing prescription or nonprescription NSAIDs (ibuprofen, naproxen, or others)
- if you take more or for a longer time than directed

Contains warning: The recommended dose of this product contains 65 mg of caffeine per tablet. Limit the use of caffeine-containing medications, such as coffee, tea, and energy drinks, while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat.

Drug Facts (continued)

Other information

- store at room temperature between 20-25°C (68 to 77°F)
- avoid excessive heat, cold and humidity
- close cap tightly after use
- read all product information before using. Keep box for important information.

Important ingredients: Ammonium Hydroxide, Black Iron Oxide, Brilliant Blue No. 1, Lactose, FD&C Red No. 40, Yellow No. 10, Benzoin Resin, FD&C Blue No. 1, Lactose, FD&C Red No. 40, Yellow No. 10, Benzoin Resin, Hydroxypropyl methylcellulose, Lactose monohydrate, Light Liquid Paraffin, Microcrystalline Cellulose, Polydioxane, Sorbitol, Stearic Acid, Talc, Titanium dioxide, Triethyl Citrate, Sorbitol, Polyethylene Glycol, Synthetic Glycerol, Sorbitol.

Questions or comments? 1-800-577-8833

Walgreens

Compare to Excedrin® Extra Strength active ingredients!

Headache Relief

ACETAMINOPHEN / ASPIRIN (NSAID) / CAFFEINE PAIN RELIEVER / PAIN RELIEVER AID

EXTRA STRENGTH GELTABS

80 GELTABS

ACTUAL SIZE

ORG0219-F

6789 12345 573742



Compare to Excedrin® Extra Strength active ingredients!

NDC 0385-1320-12

Headache Relief

ACETAMINOPHEN / ASPIRIN (NSAID) / CAFFEINE PAIN RELIEVER / PAIN RELIEVER AID

EXTRA STRENGTH GELTABS

120 GELTABS



ACTUAL SIZE

Drug Facts (continued)

Inactive ingredients
Ammonium Hydroxide, Black Iron Oxide, Brilliant Blue FCF, Corn Starch, D&C Yellow No. 10 Lake, Ethyl Cellulose, FD&C Blue No. 1 Lake, FD&C Red No. 40 Lake, Gelatin, Glycerin, Hydroxypropyl methyl Cellulose, Lactose monohydrate, Light Liquid Paraffin, Microcrystalline Cellulose, Povidone, Prepolymerized Starch, Propylene Glycol, Shellac glaze, Sorbitol Sorbitan, Stearic Acid, Talc, Titanium Dioxide, Triacetin

Questions or comments? 1-888-571-0333

CT132012 REV122018

Buy this item often?

AUTO-REORDER & SAVE

Free shipping + 20% off regular price
Learn more at Walgreens.com/AutoReorder

List No.:
Exp. Date:

Walgreens Pharmacist Recommended
Walgreens Pharmacist Verified
This product is not manufactured or distributed by Novartis Consumer Health, Inc., owner of the registered trademark Excedrin® Extra Strength.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2018 Walgreen Co.

ITEM 129612 W10058-0519-L



3 11917 20715 5



Drug Facts (continued)

With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if: you have liver disease; you are taking any medicine; you are taking any medicine for pain, fever, or headache; you have high blood pressure, heart disease, liver disease, or kidney disease; you are taking a diuretic; you are taking a blood thinner; you are taking a prescription drug for diabetes, gout, or arthritis; you are taking any other drug, or are under a doctor's care for any serious condition.

Stop use and ask a doctor if: you experience any of the following signs of stomach bleeding: vomit blood; have bloody or black stools; have stomach pain that does not get better; have dark or tarry stools; have more than 10 days of diarrhea; have more than 3 days of fever; have more than 3 days of any new symptoms occur.

These could be signs of a serious condition: If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose WARNING: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
Do not use more than directed.
Drink a full glass of water with each dose.
Use as directed: Adults 17 years and over: take 2 tablets every 6 hours; not more than 6 tablets in 24 hours.
Children under 12 years: ask a doctor.

Other Information
Store at room temperature between 20-25°C (68 to 77°F).
Avoid excessive heat, cold and humidity. Use cap tightly after use.
Read all product information before using. Keep track of important information.

Drug Facts

Temporarily relieves minor aches and pains due to: headache; cold; flu; sore throat; toothache; menstrual cramps; arthritis; muscular aches; backaches; rheumatoid arthritis; and other conditions.

Warnings
Do not use if you are allergic to aspirin, salicylates, or other drugs containing aspirin. Do not use if you are allergic to acetaminophen, acetaminophen/ aspirin, or other drugs containing acetaminophen. Do not use if you are allergic to caffeine. Do not use if you are allergic to any of the ingredients in this product. Do not use if you are allergic to any of the ingredients in this product. Do not use if you are allergic to any of the ingredients in this product. Do not use if you are allergic to any of the ingredients in this product.

Directions
Do not use more than directed. Drink a full glass of water with each dose. Use as directed: Adults 17 years and over: take 2 tablets every 6 hours; not more than 6 tablets in 24 hours. Children under 12 years: ask a doctor.



Compare to Excedrin® Extra Strength active ingredients!
NDC 0385-1320-12

Headache Relief

ACETAMINOPHEN / ASPIRIN (NSAID) / CAFFEINE PAIN RELIEVER / PAIN RELIEVER AID

EXTRA STRENGTH GELTABS

120 GELTABS



ACTUAL SIZE

ORG0219-F

20 GELTABS

EXTRA STRENGTH

ACETAMINOPHEN / ASPIRIN (NSAID) / CAFFEINE
PAIN RELIEVER / PAIN RELIEVER AID

Headache Relief

NDC 0385-1320-12



NEVER EXCEED: DO NOT USE IF IMPROPER SEAL, LOWER CAP IS DAMAGED OR MISSING

KEEP OWNER CARTON FOR COMPLETE READINGS AND PRODUCT INFORMATION

Active ingredients (in each geltab)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID*)	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses: Temporarily relieves minor aches and pains due to:

- headache ■ a cold ■ arthritis ■ muscular aches
- toothache ■ premenstrual & menstrual cramps

Warnings: **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen and Aspirin may cause severe reactions.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 geltabs in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Directions: ■ do not use more than directed

- drink a full glass of water with each dose ■ adults and children 12 years and over: take 2 geltabs every 6 hours; not more than 8 geltabs in 24 hours ■ children under 12 years: ask a doctor

Other information: ■ store at room temperature between 20-25°C (68 to 77°F) ■ avoid excessive heat, cold and humidity ■ close cap tightly after use ■ read all product information before using.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED walgreens.com ©2015 Walgreen Co.

ITEM 12562 W10206-0816-F ORG0219-F WE32208 LUC000025



ABCS

STRENGTH

TIME / ASPIRIN
PAIN RELIEVER AND

ache

1-800-333-1220

WALGREEN

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Active ingredients (in each gottab)	Purposes
Acetaminophen 250 mg.....	Pain reliever
Aspirin 250 mg (NSAID)*.....	Pain reliever
Caffeine 65 mg.....	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses Temporarily relieves minor aches and pains due to:

- headache ■ a cold ■ arthritis ■ muscular aches
- toothache ■ premenstrual & menstrual cramps

Warnings **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen and Aspirin may cause severe reactions.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 gottabs in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Directions ■ do not use more than directed

■ drink a full glass of water with each dose

■ adults and children 12 years and over: take 2 gottabs every 6 hours; not more than 8 gottabs in 24 hours

■ children under 12 years: ask a doctor

Other information

■ store at room temperature between 20-25°C (68 to 77°F)

■ avoid excessive heat, cold and humidity

■ close cap tightly after use

■ read all product information before using.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED walgreen.com ©2018 Walgreen Co

Walgreens

NDC 0383-1820-12

Headache Relief

ACETAMINOPHEN / ASPIRIN (NSAID) / CAFFEINE
PAIN RELIEVER / PAIN RELIEVER AID

EXTRA STRENGTH

GELTABS

120 GELTABS

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Active ingredients (in each gellab)	Purposes
Acetaminophen 250 mg.....	Pain reliever
Aspirin 250 mg (NSAID)*.....	Pain reliever
Caffeine 85 mg.....	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses Temporarily relieves minor aches and pains due to:

- headache ■ a cold ■ arthritis ■ muscular aches
- toothache ■ premenstrual & menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen and Aspirin may cause severe reactions.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 gellabs in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen

- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too

much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Directions

- do not use more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 tablets every 6 hours; not more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Other Information

- store at room temperature between 20-25°C (68 to 77°F)
- avoid excessive heat, cold and humidity
- close cap tightly after use
- read all product information before using.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED walgreens.com ©2018 Walgreen Co.
ITEM 129612 W10206-0819-F ORG0219-F
REV.122018 LB0363132012

ACETAMINOPHEN 250MG ASPIRIN 250MG CAFFEINE 65MG

acetaminophen 250mg aspirin 250mg caffeine 65mg tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-1320
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN (UNII: 2G86QN327L)	
D&C YELLOW NO. 10 (UNII: 35S5USQ3G)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
GLYCERIN (UNII: PDC6A3C0OX)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	

POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
STARCH, CORN (UNII: O8232NY3SJ)	
AMMONIA (UNII: 5138Q19F1X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
TALC (UNII: 7SEV7J4R1U)	
SHELLAC (UNII: 46N107B71O)	
TRIACETIN (UNII: XHX3C3X673)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

Product Characteristics

Color	green (Dark Green - White)	Score	no score
Shape	ROUND	Size	13mm
Flavor		Imprint Code	S77
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-1320-12	120 in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2019	
2	NDC:0363-1320-80	80 in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2019	
3	NDC:0363-1320-23	20 in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/24/2019	

Labeler - WALGREENS CO. (008965063)

Revised: 7/2023

WALGREENS CO.