

**MEIJER NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hbr,  
triprolidine hcl solution  
MEIJER DISTRIBUTION INC.**

-----  
**Meijer Overnight Cold & Flu**

**Drug Facts**

---

**Active ingredients (in Purposes  
each 20 mL)**

**Acetaminophen 650 mg Pain reliever/fever  
reducer**

Dextromethorphan HBr 20 Cough suppressant  
mg

Triprolidine HCl 2.5 mg Antihistamine

---

**Uses**

- temporarily relieves these common cold and flu symptoms:
  - cough
  - minor aches and pains
  - sore throat
  - headache
  - runny nose
  - sneezing
  - itching of the nose or throat
  - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

## **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

## **Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

## **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

## **When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

## **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

## **Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you

do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- **adults and children 12 years of age and older:**20 mL in dosing cup provided every 4 hours
- **children under 12 years of age:**do not use

### **Other information**

- **each 20 mL contains:**sodium 10 mg
- low sodium
- store at room temperature
- do not refrigerate

### **Inactive ingredients**

anhydrous citric acid, ascorbic acid, edetate disodium, FD&C Blue No. 1, FD&C Red No. 10, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

### **Questions or comments?**

**1-866-467-2748**

### **Principal Display Panel - 180 mL Bottle Label**

NDC 41250-869-06

Compare to Mucinex® Nightshift Cold & Flu the active ingredients\*

### **Overnight Cold & Flu**

**Acetaminophen**– Pain Reliever/Fever Reducer  
Dextromethorphan HBr – Cough Suppressant  
Triprolidine HCl – Antihistamine

### **Night Time Relief for a Better Morning**

- **Cough**
- **Fever**
- **Sore Throat**
- **Relieves runny nose**
- **Sneezing**

**For Ages 12+**

6 FL OZ (180 mL)

Tamper evident: do not use if printed seal under cap is broken or missing  
Maximum Strength per 4-hour dose.

DIST. BY:

MEIJER DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

www.meijer.com

\*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Mucinex®



## MEIJER NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, triprolidine hcl solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-869
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
<b>TRIPROLIDINE HYDROCHLORIDE</b> (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	2.5 mg in 20 mL

### Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)
<b>ASCORBIC ACID</b> (UNII: PQ6CK8PD0R)
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)
<b>PROPYL GALLATE</b> (UNII: 8D4SNN7V92)
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)
<b>WATER</b> (UNII: 059QF0K00R)
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)
<b>SORBITOL</b> (UNII: 506T60A25R)
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	FRUIT	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-869-06	180 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)	05/25/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/25/2020	

**Labeler** - MEIJER DISTRIBUTION INC. (006959555)

Revised: 10/2025

MEIJER DISTRIBUTION INC.