

LEADER NASAL DECONGESTANT- oxymetazoline hydrochloride spray
Cardinal Health 110,Inc

Leader Nasal Decongestant Spray

Drug Facts

Active Ingredient

Oxymetazoline 0.05%

Purpose

Nasal Decongestant

Uses

Temporarily relieves nasal congestion due to:

- common cold
- hay fever
- upper respiratory allergies
- temporarily relieves sinus congestion and pressure
- Shrinks swollen membranes so you can breathe more freely

Warnings

Ask a Doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- trouble urinating due to enlarged prostate gland.

When using this product

Do not use more than directed

- do not use for more than three days, Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur.
- use of this container by more than one person may spread infection.

Stop use and ask a Doctor if symptoms persist

Keep out of the reach of children

if swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Before using for the first time, remove the protective cap from the tip and prime the pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm and even stroke, and sniff deeply.
- Wipe nozzle clean after use.
- Adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- Children under 6 years of age consult a doctor.

Other information

- store between 20° - 25°C (68° to 77°F)
- retain carton for future reference on full labeling

Inactive ingredient

Benzalkonium Chloride, Disodium EDTA, Glycerin, Microcrystalline Cellulose and Carboxymethylcellulose Sodium, Polyethylene Glycol 300, Povidone, Propylene Glycol, Purified Water, Sodium Phosphate Dibasic, Sodium Phosphate Monobasic

Principal Display panel 1 FLOZ

Leader NDC 70000-0665-1

12 Hour ND Nasal Decongestant

Oxymetazoline HCL 0.05%

Nasal Decongestant

1 FLOZ (30ml)

Nasal Decongestant Spray
Maximum Strength | No Drip
 Oxymetazoline HCl 0.05% Nasal Decongestant



DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING

Drug Facts

Active ingredient Oxymetazoline hydrochloride 0.05% Nasal decongestant

Purpose
 • temporarily relieves nasal congestion due to:
 • common cold • hay fever • upper respiratory allergies
 • temporarily relieves sinus congestion and pressure
 • shrinks swollen membranes so you can breathe more freely

Warnings
Ask a doctor before use if you have
 • heart disease • high blood pressure
 • thyroid disease • diabetes
 • trouble urinating due to an enlarged prostate gland

When using this product
 • do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
 • temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur
 • use of this container by more than one person may spread infection

Stop use and ask a doctor if symptoms persist
If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
 • adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
 • children under 6 years of age: ask a doctor
To Use: Shake well before use. Before using the first time, prime metered pump by depressing pump firmly several times. Squeeze both sides of the cap and twist counterclockwise. Align the lines on the groove at the base of the pump as shown in picture. Without tilting head, insert nozzle into nostril. Depress and spray. Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use. Secure cap after use.

Child Resistant Cap (see side panel)

Drug Facts (continued)

Other information
 • Store at a controlled room temperature 20°-25°C (68°-77°F)
 • retain carton for future reference on full labeling

Inactive ingredients
 Benzalkonium Chloride, Edetate Disodium, Glycerin, Microcrystalline Cellulose and Carboxymethylcellulose Sodium, Polyethylene Glycol 300, Povidone, Propylene Glycol, Sodium Phosphate Dibasic, Sodium Phosphate Monobasic, Water (Purified)

Questions? Call 1-800-222-1087

IMPORTANT: KEEP OUTER CARTON FOR FUTURE REFERENCE ON FULL LABELING

This product is not manufactured or distributed by ©Bayer HealthCare LLC, owner of the registered trademark Afrin® Original No Drip.

All LEADER™ Brand Products Have A 100% Money Back Guarantee
 Return to place of purchase if not satisfied.



DISTRIBUTED BY CARDINAL HEALTH
 DUBLIN, OHIO 43017
 www.myleader.com 1-800-200-6315
 Essential to Care™ since 1979

©2023 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, ESSENTIAL TO CARE, LEADER, and the LEADER LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

50900



NDC 70000-0665-1

Maximum Strength | No Drip

Nasal Decongestant Spray

Oxymetazoline HCl 0.05% Nasal Decongestant

Pump Mist
12 Hour Relief

Relieves:
 Nasal Congestion
 Sinus Pressure & Congestion


COMPARE TO AFRIN® ORIGINAL NO DRIP active ingredient*

100% Money Back Guarantee

1 FL OZ (30 mL)



Directions: Squeeze both sides of the cap and twist counterclockwise. Align the lines on the groove at the base of the pump as shown in picture. Depress and spray.



Package Contains One Bottle

Actual Size

LOT

EXP

CIN 5877949 REV. 11/23





LEADER NASAL DECONGESTANT

oxymetazoline hydrochloride spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0665
Route of Administration	NASAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYMETAZOLINE HYDROCHLORIDE (UNII: K89MJ0S5VY) (OXYMETAZOLINE - UNII:8VLN5B44ZY)	OXYMETAZOLINE HYDROCHLORIDE	0.5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 300 (UNII: 5655G9Y8AQ)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
SODIUM PHOSPHATE, MONOBASIC (UNII: 3980JH2SW)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KOOR)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0665-1	1 in 1 CARTON	03/31/2024	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/31/2024	

Labeler - Cardinal Health 110,Inc (063997360)

Revised: 4/2024

Cardinal Health 110,Inc