

**MUCUS RELIEF SEVERE CONGESTION AND COUGH MAXIMUM STRENGTH-  
dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid  
TOP CARE (Topco Associates LLC)**

-----  
**Drug Facts**

**Active ingredients (in each 20 mL)**

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

**Purposes**

Cough suppressant

Expectorant

Nasal Decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep
  - nasal congestion due to a cold

**Warnings**

**Do not use**

- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

**When using this product,  
do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occur
- symptoms do not get better within 7 days or occur with a fever
- cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than 6 doses in a 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with products
- shake well before using
- mL=milliliter
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

**Other information**

- **each 20 mL contains:** sodium 17 mg
- store between 20-25°C (68°-77°F). Do not refrigerate.

**Inactive ingredients**

anhydrous citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum

**Questions or comments?**

Call toll free **1-888-423-0139**

## **Principal Display Panel**

COMPARE TO MAXIMUM STRENGTH MUCINEX® FAST-MAX® SEVERE CONGESTION & COUGH ACTIVE INGREDIENTS\*

MAXIMUM STRENGTH

### **Mucus Relief**

### **Severe Congestion and Cough**

DEXTROMETHORPHAN HBr 20 mg COUGH SUPPRESSANT

GUAIFENESIN 400 mg • EXPECTORANT

PHENYLEPHRINE HCl 10 mg • NASAL DECONGESTANT

### **MULTI-SYMPTOM RELIEF:**

- Controls Cough
- Relieves Nasal & Chest Congestion
- Thins & Loosens Mucus

FOR AGES 12 +

FL OZ (mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough.

DISTRIBUTED BY TOPCO ASSOCIATED LLC

ELK GROVE VILLAGE, IL 60007

topcare@topco.com

## **Package Label**

NDC 36800-037-06



COMPARE TO MAXIMUM STRENGTH MUCINEX® FAST-MAX® SEVERE CONGESTION & COUGH ACTIVE INGREDIENTS\*

MAXIMUM STRENGTH

# Mucus Relief Severe Congestion & Cough

DEXTROMETHORPHAN HBr 20 mg  
COUGH SUPPRESSANT  
GUAIFENESIN 400 mg • EXPECTORANT  
PHENYLEPHRINE HCl 10 mg • NASAL DECONGESTANT

### MULTI-SYMPTOM RELIEF:

- Controls Cough
- Relieves Nasal & Chest Congestion
- Thins & Loosens Mucus

PLD-E283D  
LB005241

6 FL OZ (177 mL) FOR AGES 12+

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough.



DISTRIBUTED BY TOPCO ASSOCIATES LLC  
ELK GROVE VILLAGE, IL 60007  
©TOPCO PLVA0521  
QUESTIONS? 1-888-423-0139  
topcare@topco.com  
www.topcarebrand.com  
Visit here for more information:  
<http://topbrnds.com/48zzq8>



PLD-E283D LB005242

### Drug Facts

#### Active ingredients (in each 20 mL) Purposes

Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

### Drug Facts (continued)

#### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep
  - nasal congestion due to a cold

#### Warnings

##### Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

### Drug Facts (continued)

#### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product, do not use more than directed.

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### Drug Facts (continued)

#### Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- shake well before using
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

#### Other information

- each 20 mL contains: sodium 17 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

#### Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

### Drug Facts (continued)

#### Questions or comments?

Call toll free 1-888-423-0139

PEEL CORNER FOR MORE DRUG FACTS

## TOPCARE HEALTH Maximum Strength Mucus Relief Severe Congestion & Cough

# MUCUS RELIEF SEVERE CONGESTION AND COUGH MAXIMUM STRENGTH

dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:36800-037	
<b>Route of Administration</b>	ORAL			
<b>Active Ingredient/Active Moiety</b>				
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)		GUAIFENESIN	400 mg in 20 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL	
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>			<b>Strength</b>	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
EDETATE DISODIUM (UNII: 7FLD91C86K)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYL GALLATE (UNII: 8D4SNN7V92)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)				
SORBITOL (UNII: 506T60A25R)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
XANTHAN GUM (UNII: TTV12P4NEE)				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:36800-037-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/28/2015	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
OTC Monograph Drug	M012	02/28/2015		

**Labeler** - TOP CARE (Topco Associates LLC) (006935977)