

**SINUS RELIEF PRESSURE AND PAIN- acetaminophen, guaifenesin,  
phenylephrine hcl tablet, film coated  
Kroger Company**

-----  
**Kroger 44-628**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Guaifenesin 200 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever  
Expectorant  
Nasal decongestant

***Uses***

- temporarily relieves:
  - nasal congestion
  - headache
  - sinus congestion and pressure
  - minor aches and pains
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- heart disease
- diabetes
- thyroid disease
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- liver disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

### **When using this product**

**do not exceed recommended dosage.**

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and over: take 2 caplets every 4 hours

- children under 12 years: do not use

### ***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

corn starch, crospovidone, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

### ***Questions or comments?***

**1-800-632-6900**

### ***Principal display panel***

**Kroger®**

NDC 30142-828-09

**FOR AGES 12+**

**Maximum Strength**

**Sinus Relief  
PRESSURE & PAIN**

**Acetaminophen / Pain Reliever**  
Guaifenesin / Expectorant  
Phenylephrine HCl /  
Nasal Decongestant

**RELIEVES:**  
Sinus Pressure  
& Congestion  
Headache

**Thins & Loosens Mucus**

**20**  
CAPLETS

actual  
size

**Our Pharmacist Recommended**

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER**

**UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

DISTRIBUTED BY  
THE KROGER CO.  
CINCINNATI, OHIO  
45202

QUALITY GUARANTEE  
800-632-6900  
[www.kroger.com](http://www.kroger.com)

50844 REV0419B62809

**Drug Facts (continued)**  
**Inactive ingredients** corn starch, croscollon, D & C yellow #10 aluminum lake, FD & C yellow #6 aluminum lake, magnesium stearate, talc, titanium dioxide

**Drug Facts (continued)**  
**Other information** ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN  
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
 ■ see end flap for expiration date and lot number

50844 REV0419B62809  
 PLEASE RECYCLE

**Drug Facts (continued)**  
**Warnings**  
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product  
 Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ blisters ■ rash ■ skin redness  
 If a skin reaction occurs, stop use and seek medical help right away.  
 Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 ■ you have ever had an allergic reaction to this product or any of its ingredients  
**Ask a doctor before use if you have**  
 ■ heart disease ■ diabetes ■ thyroid disease  
 ■ difficulty in urination due to enlargement of the prostate gland  
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ liver disease ■ high blood pressure  
 ■ cough that occurs with too much phlegm (mucus)

**Drug Facts (continued)**  
**Directions**  
 ■ do not take more than directed  
 ■ do not take more than 12 caplets in any 24-hour period  
 ■ adults and children 12 years and over: take 2 caplets every 4 hours  
 ■ children under 12 years: do not use

**Other information** ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN  
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
 ■ see end flap for expiration date and lot number

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

No Print / No Varnish  
 Lot no. & Exp. date

**Drug Facts**  
 NDC 30142-828-09

**Active ingredients (in each caplet)**  
 Acetaminophen 325 mg  
 Guaifenesin 200 mg  
 Phenylephrine HCl 5 mg  
 Nasal decongestant

**Purpose**  
 Pain reliever  
 Expectorant

**Uses** ■ temporarily relieves: ■ nasal congestion ■ headache ■ sinus congestion and pressure ■ minor aches and pains ■ temporarily promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Drug Facts (continued)**  
 When using this product do not exceed recommended dosage.  
 Stop use and ask a doctor if ■ nervousness, dizziness, or sleepiness occur ■ fever gets worse or lasts more than 3 days ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ redness or swelling is present ■ new symptoms occur  
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.  
 If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.  
 When using this product do not exceed recommended dosage.

85% SIZE  
 12-digit UPC-A  
 (non-suppressed)  
 For Position Only  
 0 41260 00348 6

DISTRIBUTED BY  
 THE KROGER CO.  
 CINCINNATI, OHIO  
 45202  
 QUALITY GUARANTEE  
 800-632-6900  
 www.kroger.com

B-1050-628-09-R  
 REV0419B62809

**Kroger**  
 FOR AGES 12+  
**Maximum Strength**  
**Sinus Relief**  
**PRESSURE & PAIN**  
 Acetaminophen/Pain Reliever  
 Guaifenesin/Expectorant  
 Phenylephrine HCl/  
 Nasal Decongestant



actual size

**RELIEVES:**  
 Sinus Pressure & Congestion  
 Headache  
 Thins & Loosens Mucus  
**20 CAPLETS**



**Kroger 44-628**

**SINUS RELIEF PRESSURE AND PAIN**  
 acetaminophen, guaifenesin, phenylephrine hcl tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:30142-828
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>FD&amp;C YELLOW NO. 6 ALUMINUM LAKE</b> (UNII: GYP6Z2JR6Q)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	44;628
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-828-09	2 in 1 CARTON	06/03/2019	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
OTC Monograph Drug	M012	06/03/2019	

**Labeler** - Kroger Company (006999528)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(30142-828) , pack(30142-828)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(30142-828)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(30142-828)

Revised: 4/2024

Kroger Company