

**ALLERGY MULTI-SYMPTOM- acetaminophen, chlorpheniramine maleate, phenylephrine hcl tablet, film coated**  
**L.N.K. International, Inc.**

-----  
**Quality Plus 44-455C-AMS**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Chlorpheniramine maleate 2 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever  
Antihistamine  
Nasal decongestant

***Uses***

- temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
  - headache
  - nasal congestion
  - runny nose and sneezing
  - minor aches and pains
  - sinus congestion and pressure
- temporarily relieves these additional symptoms of hay fever:
  - itchy, watery eyes
  - itching of the nose or throat
- helps clear nasal passages
- helps decongest sinus openings and passages

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- high blood pressure
- a breathing problem such as emphysema or chronic bronchitis
- heart disease
- thyroid disease
- diabetes
- liver disease
- glaucoma

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery
- drowsiness may occur

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right

away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not take more than directed**
- adults and children 12 years and over
  - take 2 caplets every 4 hours
  - swallow whole - do not crush, chew, or dissolve
  - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

### ***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

corn starch, crospovidone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

### ***Questions or comments?***

**1-800-426-9391**

### ***Principal Display Panel***

**QUALITY**

**+ PLUS**

NDC 50844-455-08

**Allergy**

**Multi-Symptom**

**Acetaminophen**

Chlorpheniramine maleate

Phenylephrine HCl

PAIN RELIEVER, ANTIHISTAMINE,  
NASAL DECONGESTANT

- Headache
- Runny Nose
- Sneezing
- Sinus Pressure
- Nasal Congestion
- Itchy, Watery Eyes

**24 Caplets**

Pseudoephedrine Free

ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS  
OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR  
SHOWS ANY SIGNS OF TAMPERING**

50844 REV0818A45508

Distributed by

**LNK INTERNATIONAL, INC.**

60 Arkay Drive

Hauppauge, NY 11788

USA



508 INC.

NDC 50844-455-08

QUALITY PLUS

# ALLERGY Multi-Symptom

Acetaminophen  
Chlorpheniramine maleate  
Phenylephrine HCl

PAIN RELIEVER, ANTIHISTAMINE,

### Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

#### Active ingredients (in each caplet)

Acetaminophen 325 mg.....Pain reliever

Chlorpheniramine maleate 2 mg.....Antihistamine

Phenylephrine HCl 5 mg.....Nasal decongestant

#### Purpose

Uses ■ temporarily relieves these symptoms of hay fever or other upper respiratory allergies:

- headache ■ nasal congestion
- runny nose and sneezing
- minor aches and pains
- sinus congestion and pressure

### Drug Facts (continued)

### Drug Facts (continued)

- temporarily relieves these additional symptoms of hay fever: ■ itchy, watery eyes
- itching of the nose or throat
- helps clear nasal passages
- helps decongest sinus openings and passages

### Warnings

- Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ blisters ■ rash ■ skin reddening
- If a skin reaction occurs, stop use and seek medical help right away.

### Do not use

- if you are now taking acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- high blood pressure

### Drug Facts (continued)

- a breathing problem such as emphysema or chronic bronchitis ■ heart disease ■ thyroid disease ■ diabetes ■ liver disease ■ glaucoma

### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery ■ drowsiness may occur

### Stop use and ask a doctor if

- nervousness, dizziness, or sleepiness occur
- pain or nasal congestion gets worse or lasts more than 7 days ■ new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- These could be signs of a serious condition.

### If pregnant or breast-feeding, ask a health

professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions


- do not take more than directed
- adults and children 12 years and over
- take 2 caplets every 4 hours
- swallow whole - do not crush, chew, or dissolve
- do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

### Drug Facts (continued)

Questions or comments? 1-800-426-9391

TAMPER EVIDENT: OPENED OR IF BLIST SHOWS ANY

B-1603-455C-08AMNSR  
REV0818A45508



Quality Plus 44-455C-AMS

## ALLERGY MULTI-SYMP TOM

acetaminophen, chlorpheniramine maleate, phenylephrine hcl tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50844-455
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>	MENTHOL	<b>Imprint Code</b>	44;455
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-455-08	2 in 1 CARTON	06/28/2005	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/28/2005	

**Labeler** - L.N.K. International, Inc. (038154464)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-455)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-455) , pack(50844-455)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50844-455)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-455)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-455)