

**COLD AND FLU RELIEF- acetaminophen, dextromethorphan hbr,  
phenylephrine hcl capsule, liquid filled  
Chain Drug Consortium, LLC**

-----

**Drug Facts**

**Active ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Purposes**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - fever
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if the user has**

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

**When using this product,**

**do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- pain, cough or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs, with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- **do not take more than directed (see overdose warning)**
- do not take more than 4 doses in 24-hour
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **When using Daytime and Nighttime products, carefully read each label to ensure correct dosing**

## Other information

- store between 15-30°C (59-86°)
- avoid excessive heat

## Inactive ingredients

FD&C yellow #6, FD&C red #40, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol white ink

## Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

## Principal Display Panel

\*COMPARE TO THE ACTIVE INGREDIENTS IN VICKS® DAYQUIL® COLD & FLU LIQUICAPS®

### Daytime, Multi-Symptom

#### Cold & Flu Relief

#### Acetaminophen 325 mg

Pain reliever/Fever reducer

#### DEXTROMETHORPHAN HBr 10 mg

Cough suppressant

#### PHENYLEPHRINE HCl 5 mg

Nasal Decongestant

Relieves: Aches • Fever • Sore Throat Cough • Nasal Congestion

Non-drowsy, Alcohol-free, Antihistamine-free

Softgels\*\*

(\*\*Liquid-filled Capsules)

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOW ANY SIGNS OR TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® DayQuil® and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

**Distributed by: Pharmacy Value Alliance, LLC**

**407 East Lancaster Avenue,**

**Wayne, PA 19087**

**Package Label**



Daytime, Multi-Symptom

# Cold & Flu Relief

**ACETAMINOPHEN 325 mg**

Pain reliever/Fever reducer

**DEXTROMETHORPHAN HBr 10 mg**

Cough suppressant

**PHENYLEPHRINE HCl 5 mg**

Nasal Decongestant

Relieves: Aches • Fever • Sore Throat  
Cough • Nasal Congestion

Non-drowsy, Alcohol-free, Antihistamine-free

**16** Softgels\*\*

Actual Size



\*COMPARE TO THE  
ACTIVE INGREDIENTS IN  
VICKS® DAYQUIL® COLD  
& FLU LIQUICAPS®

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER  
UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

No.:  
Date:

## Drug Facts (continued)

### Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

### Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

### Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

### Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**Inactive ingredients** FD&C red #40, FD&C yellow #6, gelatin, glycerin, glycol, purified water, sorbitan, sorbitol, white ink

## Drug Facts (continued)

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:  
■ more than 4,000 mg of acetaminophen in 24 hours  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure

■ trouble urinating due to an enlarged prostate gland

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

■ cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood**

thinning drug warfarin.

**When using this product, do not exceed recommended dosage.**

(\*\*Liquid-filled Capsules)

15 84

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

Distributed By: Pharmacy Value Alliance, LLC  
407 East Lancaster Avenue,  
Wayne, PA 19087



If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.



PLD-A584B FC008307 Product of China



## PREMIER VALUE Cold and Flu Relief

### COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

#### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68016-797
Route of Administration	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

#### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

#### Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	20mm

<b>Flavor</b>		<b>Imprint Code</b>		PC9
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:68016-797-16	16 in 1 CARTON	09/02/2022	09/30/2026
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>		<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug		M012	09/02/2022	09/30/2026

**Labeler** - Chain Drug Consortium, LLC (101668460)

Revised: 6/2025

Chain Drug Consortium, LLC