

**GOOD SENSE ANTIDIARRHEAL- loperamide hcl solution**  
**L. Perrigo Company**

-----  
**Perrigo Anti-Diarrheal Drug Facts**

**Active ingredient (in each 7.5 mL)**

Loperamide HCl 1 mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
children 9-11 years (60-95 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

## Other information

- **each 30 mL contains:** sodium 15 mg
- store between 20-25°C (68-77°F)

## Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

## Questions or comments?

**1-800-719-9260**

## Package/Label Principal Display Panel

Mint Flavor

See New Warning and Directions

Loperamide Hydrochloride Oral Solution, 1 mg per 7.5 mL

Anti-Diarrheal

Anti-Diarrheal Oral Solution

Controls the Symptoms of Diarrhea

Compare to active ingredient of Imodium® A-D

100% SATISFACTION GUARANTEED

4 FL OZ (120 mL)

NDC 0113-1645-26

GOODSENSE®

Mint Flavor

See New Warning and Directions

Loperamide Hydrochloride Oral Solution, 1 mg per 7.5 mL

Anti-Diarrheal

Anti-Diarrheal Oral Solution

Controls the Symptoms of Diarrhea

Compare to active ingredient of Imodium® A-D



4 FL OZ (120 mL)

Do not use if printed plastic neckband is broken or missing.

Gluten Free

Distributed By

Perrigo®

Alligan, MI 49010



0113-0645-26 1

Drug Facts

Active ingredient Purpose (in each 7.5 mL) Loperamide HCl 1 mg...Anti-diarrheal

Use controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl Heart alert: Taking more than directed can cause serious heart problems or death

Peel open to read complete directions and warnings before purchase

Drug Facts (continued)

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- fever mucus in the stool a history of liver disease a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse diarrhea lasts for more than 2 days you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea find right dose on chart. If possible, use weight to dose; otherwise use age. shake well before using

ADHESIVE AREA NO COPY

ADHESIVE AREA NO COPY

Drug Facts (continued)

- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device. mL = milliliter

Table with 2 columns: Age/Weight Group and Dosage. Rows include adults and children 12 years and over, children 9-11 years (60-95 lbs), children 6-8 years (48-59 lbs), children 2-5 years (34 to 47 lbs), and children under 2 years (up to 33 lbs).

Other information

- each 30 mL contains: sodium 15 mg store between 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

Questions or comments? 1-800-719-9260

64526 C2 F4

# GOOD SENSE ANTIDIARRHEAL

loperamide hcl solution

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0113-1645
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	1 mg in 7.5 mL

## Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XANTHAN GUM (UNII: TTV12P4NEE)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

## Product Characteristics

Color	GREEN (opaque, viscous)	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-1645-26	120 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/21/2018	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091292	05/21/2018	

**Labeler** - L. Perrigo Company (006013346)

Revised: 1/2020

L. Perrigo Company