

IBUPROFEN- ibuprofen capsule, liquid filled
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*

(present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - trouble breathing
 - leg swelling
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
 - take 1 capsule every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 capsule, 2 capsules may be used
 - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- **each capsule contains:** potassium 20 mg
- store at 20 - 25°C (68 - 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.
- read all warnings and directions before use. Keep carton.
- swallow whole, do not crush, chew, or dissolve

Inactive ingredients

FD&C blue #1, FD&C yellow#6, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

Questions or comments?

Call toll free **1-888-309-9030**

Principal Display Panel

Compare to the active ingredient of Advil® Liqui-Gel®†

Ibuprofen

Solubilized Ibuprofen Capsules, 200 mg

Pain Reliever / Fever Reducer **(NSAID)**

Softgels**

**Liquid Filled Capsules

†This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

TAMPER EVIDENT: DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

PACKAGED BY OLD EAST MAIN CO.

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072 USA

Product Label

DG™ health

Compare to the active ingredient of Advil® Liqui-Gels^{®†}

Ibuprofen

Solubilized Ibuprofen Capsules, 200 mg
Pain Reliever/Fever Reducer (NSAID)

200
mg

40 Softgels**

(*Liquid-Filled Capsules)

Actual Size



<p>Drug Facts</p> <p>Active ingredient (in each capsule) Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)* (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug</p> <p>Purpose Pain reliever/fever reducer</p>	<p>Uses</p> <ul style="list-style-type: none"> temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> muscular aches minor pain of arthritis toothache backache menstrual cramps temporarily reduces fever
<p>Warnings</p> <p>Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> hives facial swelling skin redness shock skin redness asthma (wheezing) rash blisters <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"> are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Do not use</p> <ul style="list-style-type: none"> if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery 	

<p>Drug Facts (continued)</p> <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke you are taking a diuretic <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> under a doctor's care for any serious condition taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug <p>When using this product</p> <ul style="list-style-type: none"> take with food or milk, if stomach upset occurs <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> feel faint vomit blood have bloody or black stools have stomach pain that does not get better chest pain slurred speech leg swelling trouble breathing weakness in one part or side of body pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present in the painful area any new symptoms appear <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>

<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than directed the smallest effective dose should be used adults and children 12 years and over: <ul style="list-style-type: none"> take 1 capsule every 4 to 6 hours while symptoms persist if pain or fever does not respond to 1 capsule, 2 capsules may be used do not exceed 6 capsules in 24 hours, unless directed by a doctor children under 12 years: ask a doctor <p>Other information</p> <ul style="list-style-type: none"> each capsule contains: potassium 20 mg store at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F). Protect from light. read all warnings and directions before use. Keep carton. swallow whole; do not crush, chew, or dissolve <p>Inactive ingredients FD&C blue no. 1, FD&C yellow no. 6, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan</p> <p>Questions or comments? Call 1-888-309-9030</p>

This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

DISTRIBUTED BY OLD EAST MAIN CO.
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072

100%
Satisfaction
Guaranteed!
(888) 309-9030

Exp. Date:
Lot No.:

P.D.-AS298
F007039



TAMPER EVIDENT: DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING. READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

DOLLAR GENERAL HEALTH Ibuprofen Softgels

IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-750
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	AT146
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-750-12	12 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2018	01/31/2025
2	NDC:55910-750-80	1 in 1 BOX	08/31/2018	01/31/2025
2		80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:55910-750-02	1 in 1 BOX	08/31/2018	01/31/2025
3		40 in 1 BOTTLE, PLASTIC; Type 0: Not a		

3		Combination Product		
4	NDC:55910-750-20	1 in 1 BOX	08/31/2018	01/31/2025
4		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
5	NDC:55910-750-40	1 in 1 BOX	08/31/2018	01/31/2025
5		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206999	08/31/2018	01/31/2025

Labeler - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 10/2022

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)