

SEVERE COLD AND FLU- acetaminophen, diphenhydramine hydrochloride, phenylephrine hydrochloride powder, for solution
Walgreen Company

Walgreen Co. Severe Cold & Flu Drug Facts

Active ingredients (in each packet)

Acetaminophen 650 mg

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Antihistamine/cough suppressant

Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
- minor aches and pains
- minor sore throat pain
- headache
- nasal and sinus congestion
- runny nose
- sneezing
- itchy nose or throat
- itchy, watery eyes due to hay fever
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not exceed recommended dosage**
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed (see overdose warning)**
- take every 4 hours, while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

Other information

- **each packet contains:** potassium 10 mg and sodium 25 mg
- **phenylketonurics:** contains phenylalanine 13 mg per packet
- store at 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients

acesulfame potassium, anhydrous citric acid, aspartame, colloidal silicon dioxide, D&C yellow #10, FD&C blue #1, FD&C red #40, flavors, maltodextrin, pregelatinized starch, sodium citrate, sucrose, tribasic calcium phosphate

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Walgreens

Compare to the active ingredients in Theraflu® Severe Cold & Cough

WALGREENS PHARMACIST RECOMMENDED

NIGHTTIME

Severe Cold & Flu

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER

DIPHENHYDRAMINE HCl / ANTIHISTAMINE / COUGH SUPPRESSANT

PHENYLEPHRINE HCl / NASAL DECONGESTANT

● Relieves nasal congestion, cough, body ache, sore throat pain, runny nose, sneezing, headache & fever

6 PACKETS

Honey Lemon

infused with white tea flavors

NDC 0363-1357-01

Walgreens



Compare to the active ingredients
in Theraflu® Severe Cold & Cough^{††}

NIGHTTIME

Severe Cold & Flu

**ACETAMINOPHEN / PAIN RELIEVER /
FEVER REDUCER
DIPHENHYDRAMINE HCl / ANTIHISTAMINE /
COUGH SUPPRESSANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT**

- Relieves nasal congestion, cough,
body ache, sore throat pain,
runny nose, sneezing,
headache & fever



6

PACKETS

Honey Lemon
infused with white tea flavors



[†]Our pharmacists recommend the
Walgreens brand. We invite you to
compare to national brands.

^{††}This product is not manufactured or
distributed by Novartis Consumer Health
S.A., owner of the registered trademark
Theraflu®.

W3ORG0122-F



CONVENIENT RECLOSEING TAB

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

Drug Facts

Active ingredients (in each packet)

Purposes

Acetaminophen 650 mg.....	Pain reliever/fever reducer
Diphenhydramine HCl 25 mg.....	Antihistamine/cough suppressant
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - minor sore throat pain
 - headache
 - nasal and sinus congestion
 - runny nose
 - sneezing
 - itchy nose or throat
 - itchy, watery eyes due to hay fever
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)
- take every 4 hours, while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 10 mg and sodium 25 mg
- phenylketonurics: contains phenylalanine 13 mg per packet
- store at 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients

acesulfame potassium, anhydrous citric acid, aspartame, colloidal silicon dioxide, D&C yellow #10, FD&C blue #1, FD&C red #40, flavors, maltodextrin, pregelatinized starch, sodium citrate, sucrose, tribasic calcium phosphate

Questions or comments?

1-800-719-9260

Gluten Free

OPEN OTHER END

DISTRIBUTED BY: WALGREEN CO.
 200 WILMOT RD., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED
 walgreens.com ©2021 Walgreen Co.
 MADE IN MEXICO

ITEM 538582 W10106-0522-L



**CODE
AREA**

96491 94 C5

SEVERE COLD AND FLU

acetaminophen, diphenhydramine hydrochloride, phenylephrine hydrochloride powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-1357
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
ASPARTAME (UNII: Z0H242BBR1)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SUCROSE (UNII: C151H8M554)	
TRIBASIC CALCIUM PHOSPHATE (UNII: 91D9GV0Z28)	

Product Characteristics

Color	WHITE (mixture of white, light yellow-orange particles) , ORANGE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
----------	------------------	----------------------------	-----------------------------	---------------------------

1	NDC:0363-1357-91	6 in 1 CARTON; Type 0: Not a Combination Product	05/31/2023	02/01/2025
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		05/31/2023	02/01/2025

Labeler - Walgreen Company (008965063)

Revised: 7/2024

Walgreen Company