

FAMOTIDINE- famotidine tablet, film coated FAMILY DOLLAR

Famotidine Tablets

Drug Facts

Active ingredient (in each tablet)

Famotidine, USP 10 mg and 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert

Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet (of 10 mg) with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - to **prevent** symptoms, swallow 1 tablet (of 20 mg) with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - **before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°C to 25°C (68°F to 77°F)
- protect from moisture

Inactive ingredients

Microcrystalline Cellulose, Pre-gelatinized Starch, Sodium Starch glycolate, Hydroxypropyl Cellulose, Magnesium stearate, Hypromellose, Titanium dioxide, Triacetin, Talc, Macrogol

Questions or comments?

Call us at **1 (888) 721-7115 (toll-free)**

Distributed by: MIDWOOD BRANDS, LLC

500 Volvo Pkwy, Chesapeake, Va 23320 USA

Product of India

M.L. No.: 02/SKL/AP/2015/F/R

Principal Display Panel

NDC 55319-408-01

Famotidine Tablets, USP 10mg

60 Tablets

FAMILY
Wellness™

ORIGINAL STRENGTH
FAMOTIDINE

TABLETS, USP 10 mg
Acid Reducer

JUST ONE TABLET!

Prevents & relieves heartburn
due to acid indigestion

60 TABLETS NDC 55319-408-01

Drug Facts

Active ingredient (in each tablet) Purpose
Famotidine, USP 10 mg.....Acid reducer

Uses
■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers
Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
■ with other acid reducers
Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating, or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn

■ unexplained weight loss ■ nausea or vomiting
■ stomach pain ■ kidney disease
Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
■ adults and children 12 years and over:
■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn
■ do not use more than 2 tablets in 24 hours
■ children under 12 years: ask a doctor

TAMPER EVIDENT: DO NOT USE IF PRINTED FOIL SEAL ON BOTTLE IS BROKEN OR MISSING.

Other information
■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20°C to 25°C (68°F to 77°F) ■ protect from moisture

Inactive ingredients
microcrystalline cellulose, pre-gelatinized starch, sodium starch glycolate, hydroxypropyl cellulose, magnesium stearate, hypromellose, titanium dioxide, triacetin, talc, macrogol

Questions or Comments?
In the U.S., call weekdays 9 AM to 6 PM at 1 (888) 721-7115.
DISTRIBUTED BY: MIDWOOD BRANDS, LLC
500 VOLVO PKWY, CHESAPEAKE, VA 23320 USA **PRODUCT OF INDIA**
NOT 100% SATISFIED? Return within 30 days to the store of purchase for a refund (with receipt) or exchange.
Mfg.Lic.No.: 02/SKL/AP/2015/F/R 05/25 SKU 997772

LOT

EXP

800340

Principal Display Panel

NDC 55319-409-03

Famotidine Tablets, USP 20mg

100 Tablets

FAMILY
Wellness™

MAXIMUM STRENGTH
FAMOTIDINE

TABLETS, USP 20 mg
Acid Reducer

JUST ONE TABLET!

Prevents & relieves heartburn
due to acid indigestion

100 TABLETS NDC 55319-409-03

Drug Facts

Active ingredient (in each tablet) Purpose
Famotidine, USP 20 mg.....Acid reducer

Uses
■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers
Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
■ with other acid reducers
Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating, or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn

■ unexplained weight loss ■ nausea or vomiting
■ stomach pain ■ kidney disease
Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
■ adults and children 12 years and over:
■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
■ do not use more than 2 tablets in 24 hours
■ children under 12 years: ask a doctor

TAMPER EVIDENT: DO NOT USE IF PRINTED FOIL SEAL ON BOTTLE IS BROKEN OR MISSING.

Other information
■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20°C to 25°C (68°F to 77°F) ■ protect from moisture

Inactive ingredients
microcrystalline cellulose, pre-gelatinized starch, sodium starch glycolate, hydroxypropyl cellulose, magnesium stearate, hypromellose, titanium dioxide, triacetin, talc, macrogol

Questions or Comments?
In the U.S., call weekdays 9 AM to 6 PM at 1 (888) 721-7115.
DISTRIBUTED BY: MIDWOOD BRANDS, LLC
500 VOLVO PKWY, CHESAPEAKE, VA 23320 USA **PRODUCT OF INDIA**
NOT 100% SATISFIED? Return within 30 days to the store of purchase for a refund (with receipt) or exchange.
Mfg.Lic.No.: 02/SKL/AP/2015/F/R 05/25 SKU 997790

LOT

EXP

800343

FAMOTIDINE

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-408
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg

Inactive Ingredients	
Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
HYDROXYPROPYL CELLULOSE (110000 WAMW) (UNII: 5Y0974F5PW)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HFI6D95)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics			
Color	WHITE (white to off-white)	Score	no score
Shape	ROUND	Size	5mm
Flavor		Imprint Code	V;21
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-408-01	1 in 1 CARTON	09/01/2025	
1		60 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215822	09/01/2025	

FAMOTIDINE
famotidine tablet, film coated

Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	NDC:55319-409
Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)			FAMOTIDINE	20 mg
Inactive Ingredients				
Ingredient Name				Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
STARCH, CORN (UNII: O8232NY3SJ)				
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)				
HYDROXYPROPYL CELLULOSE (110000 WAMW) (UNII: 5Y0974F5PW)				
MAGNESIUM STEARATE (UNII: 70097M6I3O)				
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
TRIACETIN (UNII: XHX3C3X673)				
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HFI6D95)				
TALC (UNII: 7SEV7J4R1U)				
Product Characteristics				
Color	WHITE (white to off-white)		Score	no score
Shape	ROUND		Size	6mm
Flavor			Imprint Code	V;15
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-409-01	1 in 1 CARTON	09/01/2025	
1		25 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55319-409-02	1 in 1 CARTON	09/01/2025	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:55319-409-03	1 in 1 CARTON	09/01/2025	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215822	09/01/2025	

Labeler - FAMILY DOLLAR (024472631)

Establishment

Name	Address	ID/FEI	Business Operations
VKT Pharma Private Limited		871408062	MANUFACTURE(55319-408, 55319-409)

Revised: 6/2025

FAMILY DOLLAR