

TOOTHACHE AND GUM RELIEF CUSHIONS- benzocaine dressing
Rite Aid

5820616 RA Tooth Cushions

Benzocaine 10% . . . Oral pain reliever

Menthol 0.25% . . . Oral pain reliever

for the temporary relief of pain due to toothaches

Methemoglobinemia warning: use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develop: * pale, gray, or blue colored skin (cyanosis) * headache * rapid heart rate * shortness of breath * dizziness or lightheadedness * fatigue or lack of energy

Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics

Do not use * more than directed * for more than 7 days unless told to do so by a dentist or doctor * for teething * in children under 2 years of age

Stop use and ask a doctor if * swelling, rash or fever develops * irritation, pain or redness persists or worsens * symptoms do not improve in 7 days * allergic reaction occurs

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

carefully remove gel cushion from blister. Gently mold with fingers until soft. Apply to the affected tooth/gum area and gently press for 5-10 seconds to mold for a comfortable fit. Gel cushion will stay in place and dissolve slowly in mouth. Adults and children 12 years of age and older: use up to 3 times daily or as directed by a dentist or doctor. Children under 12 years of age: ask a dentist or doctor. Children under 2 years of age: do not use

Do not use if blister card is broken. This preparation is intended for use in cases of toothache, only as a temporary expedient until a dentist can be consulted. Do not use continuously. These toothache cushions are designed to completely dissolve in place. Store at room temperature 15-30°C (59-86°F)

cellulose gum, flavor, polyethylene glycol



TOOTHACHE AND GUM RELIEF CUSHIONS

benzocaine dressing

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-1616
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	0.25 g in 100 g
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	10 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	

Product Characteristics

Color	white (Off white to beige)	Score	
Shape		Size	
Flavor	MINT (Mint with menthol)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-1616-9	8 in 1 CARTON	06/26/2020	
1		0.5 g in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	06/26/2020	

Labeler - Rite Aid (014578892)

Registrant - Lornamead (119559733)

Establishment

Name	Address	ID/FEI	Business Operations
Lornamead Products		119559733	manufacture(11822-1616) , pack(11822-1616)

Revised: 11/2025

Rite Aid