

**LORATADINE- loratadine solution**  
**ATLANTIC BIOLOGICALS CORP.**

-----  
**Children's Loratadine Oral Solution**

**Loratadine Oral Solution - Allergy**

Active Ingredient: Loratadine 5 mg (in each 5 mL(teaspoonful)(TSP))

Purpose: Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

**Warnings**

**Do not use if you have** ever had allergic reaction to this product or any of it's ingredients

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**When using this product** do not take more than directed. Taking more than directed may cause drowsiness.

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

---

adults and children 6 years and over	2 teaspoonfuls (TSP) daily; do not take more than 2 teaspoonfuls (TSP) in 24 hours
children 2 to under 6 years of age	1 teaspoonful (TSP) daily; do not take more than 1 teaspoonful (TSP) in 24 hours
children under 2 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

---

**Other information**

- **Each teaspoonful contains:** sodium 1 mg

- Safety Sealed. Do not use if imprinted seal around cap is broken or missing
- Store between 20° to 25°C (68° to 77°F)

### **Inactive ingredients**

Butylated hydroxyanisole, citric acid, glycerin, grape flavor, propylene glycol, purified water, sodium benzoate, sucralose.

### **Questions**

#### **DISTRIBUTED BY:**

ATLANTIC BIOLOGICALS CORP.

MIAMI, FL 33179

### **Loratadine Oral Solution - Hives Relief**

Active Ingredient: Loratadine 5 mg (in each 5 mL(teaspoonful(tsp)))

Purpose: Antihistamine

### **Uses**

- relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

### **Warnings**

**Severe Allergy Warning:** Get emergency help **immediately** if you have hives along with any of the following symptoms:

- trouble swallowing
- dizziness or loss of consciousness
- swelling of tongue
- swelling in or around mouth
- trouble speaking
- drooling
- wheezing or problems breathing

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

**Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for “anaphylaxis” or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

### **Do not use**

- to **prevent** hives from any known cause such as:
  - foods
  - insects stings

- medicines
- latex or rubber gloves

because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.

- if you have ever had an allergic reaction to this product (loratadine) or any of its ingredients

**Ask a doctor before use if you have**

- liver or kidney disease. Your doctor should determine if you need a different dose.
- hives that are an unusual color, look bruised or blistered
- hives that do not itch

**When using this product** do not take more than directed. Taking more than directed may cause drowsiness.

**Stop use and ask a doctor if**

- **an allergic reaction to this product occurs. Get medical help right away**
- symptoms don't improve after 3 days of treatment
- the hives have lasted more than 6 weeks

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

---

adults and children 6 years and over	2 teaspoonfuls (tsp) daily; do not take more than 2 teaspoonfuls (tsp) in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

---

**Other information**

- Safety sealed. Do not use if imprinted seal around cap is broken or missing
- Store between 20° to 25°C (68° to 77°F)

**Inactive ingredients**

Butylated hydroxyanisole, citric acid, glycerin, grape flavor, propylene glycol, purified water, sodium benzoate, sucralose.

**Questions**

DISTRIBUTED BY:

ATLANTIC BIOLOGICALS CORP.

MIAMI, FL 33179

**Patient Leaflet**

**Once Daily / Non-Drowsy\***

**Loratadine Oral Solution**

**Hives Relief**

**5 mg per 5 mL**

**Antihistamine**

**What you should know about:**

**Non- Drowsy\***

**Loratadine Oral Solution – Hives Relief**

**Original Prescription Strength – 24 Hour Product**

**Relief of Itching Due to Hives**

(Please read all of this information before taking Loratadine Oral Solution – Hives Relief Product. Save this leaflet for future reference.)

**What is Loratadine Oral Solution - Hives Relief product?**

Loratadine Oral Solution - Hives Relief product contain loratadine, a non-drowsy \*antihistamine.

---

**Loratadine Oral Solution – Hives Relief**

**Original prescription strength medicine to relieve itching due to hives.**

**What are hives and what do they look like?**

Hives are itchy raised red or pink bumps, blotches or streaky areas of various sizes on the surface of the skin. They can come and go.

**Do not use this product if your hives look bruised, have an unusual color, or look blistered. These may be signs of a more serious problem. If you have these signs, ask a Doctor before taking Loratadine Oral Solution -Hives Relief product.**

**What can cause hives?** Many things, including food, drugs (medicines), insect stings, stress, exercise or temperature can cause hives. A doctor may be able to help you find the cause of your hives. When you know the cause, avoid it.

**Do not use this product to prevent hives or allergic skin reactions known to be caused by food, insect stings, medicines, latex or rubber gloves, or any other causes. Many medicines can cause hives.**

**How long do hives last?** Hives can last from a few hours to a few weeks, and often come and go. See a doctor if your hives have lasted more than 6 weeks.

**Are hives very serious?**

Hives are very common and they are usually not serious. **But in rare cases, they can be the first sign of a life threatening allergic reaction** called anaphylaxis. A sign that you have this serious reaction would be that you have hives along with any of the following symptoms:

- Dizziness or feeling faint
- Trouble swallowing
- Wheezing or problems breathing
- Trouble speaking
- Hives or swelling in or around the mouth
- Drooling

**Hives AND any of these symptoms means call 911 or Get Immediate Medical Help**

## **Can I use Loratadine Oral Solution -Hives Relief product instead of an epinephrine injection for my allergic reaction?**

**Never use Loratadine Oral Solution – Hives Relief product as a substitute for epinephrine.** If your doctor has prescribed an epinephrine injector for your severe allergic reactions, your doctor is concerned that your hives might lead to a severe allergic reaction (anaphylaxis) or to asthma-like symptoms that could occur with your hives. Loratadine Oral Solution – Hives Relief product is not a substitute for epinephrine. If you have been prescribed an epinephrine injector, carry it with you at all times.

### **Who can take Loratadine Oral Solution – Hives Relief Product?**

People who have hives (urticaria) and are 6 years of age or older can take Loratadine Oral Solution – Hives Relief product. (See label Directions for proper dose.)

### **Who should not take Loratadine Oral Solution – Hives Relief Product?**

- People who have ever had an allergic reaction to these products or any of their ingredients.
- People who are taking any other products containing loratadine
- People with liver or kidney disease unless directed by a doctor
- Children under the age of 6 years.

### **What if I know the cause of my hives and I want to take Loratadine Oral Solution – Hives Relief Product to prevent the hives from occurring?**

Loratadine Oral Solution – Hives Relief product should never be used to prevent hives or allergic skin reactions from any known cause. Such causes can include but are not limited to:

- Food
- Insect Stings
- Medicines
- Latex or rubber gloves.

If you have ever had an allergic reaction to any of these causes, you should see a doctor.

### **When should you contact a doctor?**

**Contact a doctor before taking Loratadine Oral Solution – Hives Relief product** if you have:

- Hives that have an unusual color or that look bruised or blistered
- Hives that have lasted longer than 6 weeks

If you are pregnant or breast feeding, you should ask a doctor or health professional before use.

**Contact a doctor while taking Loratadine Oral Solution – Hives Relief Product** if you (or your child) have:

- A rash or itching that does not get better after taking Loratadine Oral Solution – Hives Relief product for 3 days
- Hives that keep coming back.

### **How long can Loratadine Oral Solution – Hives Relief product be taken to relieve the itching?**

Loratadine Oral Solution – Hives Relief product can be taken during the entire time that you have itching due to hives (see Directions). Don't forget, if the hives last longer than 6 weeks, contact your doctor.

### **How should Loratadine Oral Solution – Hives Relief product be taken?**

- Adults and children 6 years and older should take 2 teaspoonfuls of Loratadine Oral Solution (10 mg) daily.

- Do not take more than one dose in 24 hour period.
- If you have liver or kidney disease, ask a doctor.

**What else can I do to make the itching and rash feel better?**

Wear loose clothing to help reduce irritation. Keeping the temperature cool can help with the itching.

**Other important information:**

- Allergy Alert: Do not use if you are allergic to loratadine, any of its ingredients, or other antihistamines.
- If you have been prescribed an epinephrine injector, you should carry it with you at all times.
- Keep out of reach of children.
- In case of an overdose, get medical help or contact a Poison Control Center right away.

\* When taken as directed. See drug facts panel on carton.

NDC 17856-0557-1

**17856-0557-01**



Rev.01/19

**LORATADINE ORAL SOLUTION  
5 MG/5 ML ALCOHOL  
FREE/SUGAR FREE/DYE FREE**

See package insert for indications and dosage schedule

Store between 20° to 25°C (68° to 77°F). Each teaspoonful (TSP) contains: sodium 1 mg  
GRAPE FLAVOR  
KEEP OUT OF THE REACH OF CHILDREN


**17856-0557-01  
LORATADINE**

Dosage: 5 ML

Qty: 72 Cups



GTIN: 1785605571  
S/N: 00000000000000  
Exp: 02/07/20  
Lot: 000000



17856055701

Packaged by: Unit Dose Solutions  
Morrisville, NC 27560

Distributed by: AtlanticBiologicals  
Corp, Miami FL 33179

Call to Reorder: 800.509.7592

**LORATADINE**

loratadine solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:17856-0557(NDC:54838-558)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
LORATADINE (UNII: 7AJ03BO7QN) (LORATADINE - UNII:7AJ03BO7QN)	LORATADINE	5 mg in 5 mL

### Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
WATER (UNII: 059QF0K00R)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE (grape flavor)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17856-0557-1	72 in 1 BOX, UNIT-DOSE	01/22/2021	
1		5 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA077421	02/13/2019	

**Labeler** - ATLANTIC BIOLOGICALS CORP. (047437707)

**Establishment**

Name	Address	ID/FEI	Business Operations
ATLANTIC BIOLOGICALS CORP.		047437707	repack(17856-0557) , label(17856-0557)

Revised: 1/2021

ATLANTIC BIOLOGICALS CORP.