

**ADULT TUSSIN COUGH AND CHEST CONGESTION DM- dextromethorphan hbr,  
guaifenesin liquid  
Fred's Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredients (in each 10 mL)**

Dextromethorphan HBr 20 mg

Guaifenesin 200 mg

**Purposes**

Cough suppressant

Expectorant

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

**Warnings**

**Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Stop use and ask a doctor if**

cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than 6 doses in any 24-hour period

- measure only with the dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL= milliliter
- this adult product is not intended for use in children under 12 years of age
- adults and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

### **Other information**

- store between 20-25°C(68-77°F). Do not refrigerate

### **Inactive ingredients**

anhydrous citric acid, FD&C red 40, flavor, glucose, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate

### **Questions or comments?**

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

### **Principal Display Panel**

Compare To The Active Ingredients in Pobitussin® Peak Cold Cpough-+Chest Congestion DM\*

#### **Adult**

#### **Tussin Cough + Chest Congestion DM**

DEXTROMETHORPHAN HBr 20 mg (Cough Suppressant)

GUAIFENESIN 200 mg (Expectorant)

Relieves:

- Cough
- Mucus

NON-DROWSY

ALCOHOL-FREE

FL OZ (mL)

DOSAGE CUP INCLUDED

FOR AGES 12 & OVER

\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributors of Robitussin® Peak Cold Cough + Chest Congestion DM.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

DISTRIBUTED BY: fred's, Inc.

4300 NEW GETWELL RD,

MEMPHIS, TN 38118

[www.fredsinc.com](http://www.fredsinc.com)

### **Package Label**

**Drug Facts (continued)**

**Other information**  
 ■ store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, FD&C red #40, flavor, glucose, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate

**Questions or comments?**  
 Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Robitussin® Peak Cold Cough+Chest Congestion DM.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**PARENTS:**  
 Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**100%** satisfaction guaranteed  
 Questions or comments:  
 1-866-331-FRED (3733)

DISTRIBUTED BY: fred's, Inc.  
 4300 NEW GETWELL RD,  
 MEMPHIS, TN 38118  
[www.fredsinc.com](http://www.fredsinc.com)

**fred's Pharmacy** NDC 55315-385-04

Compare to the active ingredients in Robitussin® Peak Cold Cough+Chest Congestion DM\*

**ADULT**

# Tussin

## COUGH + CHEST CONGESTION DM

DEXTROMETHORPHAN HBr 20mg (Cough Suppressant)  
GUAIFENESIN 200mg (Expectorant)

**RELIEVES:**

- COUGH
- MUCUS

**NON-DROWSY ALCOHOL-FREE**

Dosing Cup Included

**4 FL OZ (118 mL)**

**FOR AGES 12 & OVER**

**fred's Pharmacy** NDC 55315-385-04

Compare to the active ingredients in Robitussin® Peak Cold Cough+Chest Congestion DM\*

**ADULT**

# Tussin

## COUGH + CHEST CONGESTION DM

DEXTROMETHORPHAN HBr 20mg (Cough Suppressant)  
GUAIFENESIN 200mg (Expectorant)

**RELIEVES:**

- COUGH
- MUCUS

**NON-DROWSY ALCOHOL-FREE**

Dosing Cup Included

**4 FL OZ (118 mL)**

**FOR AGES 12 & OVER**

**Drug Facts**

Active ingredients (in each 10 mL)	Purposes
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

**Warnings**

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Stop use and ask a doctor if** cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.**  
 In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adults and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use



PLD-K325C FC004390

Lot No.:  
 Exp. Date:

**FRED'S PHARMACY Adult Tussin Cough + Chest Congestion DM**

**ADULT TUSSIN COUGH AND CHEST CONGESTION DM**

dextromethorphan hbr, guaifenesin liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55315-385
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL

## Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
DEXTROSE (UNII: IY9XDZ35W2)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
MENTHOL (UNII: L7T10EIP3A)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55315-385-08	1 in 1 BOX	06/30/2015	
1		237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:55315-385-04	1 in 1 BOX	06/30/2015	
2		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	06/30/2015	

**Labeler** - Freds Inc (005866116)