

DOUBLE DIP- iodine teat dip solution

Eastern Crown Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

ECI Double Dip Iodine Teat Dip

DIRECTIONS FOR USE

Use at full strength. Do not dilute.

Before starting or continuing to use teat dip, consult your veterinarian if cow's teats are sore or chapped.

ECI Double Dip is manufactured for double use as both a before and after milking dip.

Before Milking:

- Remove excess organic debris from teats and forestrip. If water is used, use minimal amount. Thoroughly dry with a single service towel.
- Check foremilk for abnormalities.
- Dip or spray entire cow's teats with ECI DOUBLE DIP.
- Allow 15 to 30 seconds contact time.
- Dry thoroughly, with clean, single service paper towel.
- Attach milking machine

After Milking:

- Immediately after milking, dip or spray each entire teat with ECI DOUBLE DIP. Allow to dry. **Do not wipe.** Before next milking, wash cow's udder and teats with potable water and dry with a single service paper towel.
- Appropriately clean teat dipper after each milking.

When cow is being dried off, the teats should be dipped once a day at least 3 or 4 days after the last milking.

When freshening cows, begin dipping teats with Double Dip twice a day for 3-4 days before calving.

DIP SOLUTION SHOULD BE CHANGED IF IT BECOMES VISIBLY DIRTY, IF SEDIMENT IS PRESENT, OR IF THE AMBER COLOR FADES NOTICEABLY.

Double Dip may aid in controlling the spread of organisms that cause mastitis.

Do not use for cleaning or sanitizing milking equipment.

STORAGE AND DISPOSAL:

DO NOT ALLOW TO FREEZE.

Store in cool dry area. Shake well before using. Keep container tightly closed when not in use. Do not reuse empty container. Rinse container before disposing in a safe manner. Avoid storing near products containing chlorine.

NFPA = 100

CAUTION!

Can be severely irritating to eyes. Can be irritating to skin and harmful if swallowed.

First Aid

ORAL INGESTION: If swallowed, dilute by giving several glasses of water or milk. Do not give anything by mouth to an unconscious person. Do not induce vomiting. If vomiting does occur, repeat giving several glasses of water or milk and consult a physician.

EYE CONTACT: Flush eyes with plenty of clean running water for at least 15 minutes. If irritation occurs and persists, get medical attention.

SKIN CONTACT: Wash thoroughly. If irritation occurs and persists, get medical attention.

INHALATION: Remove victim to fresh air.

CAUTION!
Can be severely irritating to eyes. Can be irritating to skin and harmful if swallowed.

First Aid
ORAL INGESTION: If swallowed, dilute by giving several glasses of water or milk. Do not give anything by mouth to an unconscious person. Do not induce vomiting. If vomiting does occur, repeat giving several glasses of water or milk and consult a physician.
EYE CONTACT: Flush eyes with plenty of clean running water for at least 15 minutes. If irritation occurs and persists, get medical attention.
SKIN CONTACT: Wash thoroughly. If irritation occurs and persists, get medical attention.
INHALATION: Remove victim to fresh air.

STORAGE AND DISPOSAL:
DO NOT ALLOW TO FREEZE.
Store in cool dry area. Shake well before using. Keep container tightly closed when not in use. Do not reuse empty container. Rinse container before disposing in a safe manner. Avoid storing near products containing chlorine.

NFPA=100

NET CONTENT: Gallons

LOT NO.:

EXPIRATION DATE:

ECI

NPE Free

DOUBLE DIP
IODINE TEAT DIP
NDC 060357-1010

ACTIVE INGREDIENT: Iodine (titratable): 1.0%
OTHER INGREDIENTS: 99.0% Contains Glycerin and Lanolin.
Contains No Nonylphenoxy Poly Ethanol (NPE)

KEEP OUT OF REACH OF CHILDREN.
CAUTION
Harmful if Swallowed • Eye Irritant
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS.

Directions for Use:
Use at full strength. Do not dilute.
Before starting or continuing to use teat dip, consult your veterinarian if cow's teats are sore or chapped. ECI Double Dip is manufactured for double use as both a before and after milking dip.

Before Milking:

- Remove excess organic debris from teats and forestrip. If water is used, use minimal amount. Thoroughly dry with a single service towel.
- Check foremilk for abnormalities.
- Dip or spray entire cow's teats with ECI DOUBLE DIP.
- Allow 15 to 30 seconds contact time.
- Dry thoroughly, with clean, single service paper towel.
- Attach milking machine.

After Milking:

- Immediately after milking, dip or spray each entire teat with ECI DOUBLE DIP. Allow to dry. **Do not wipe.** Before next milking, wash cow's udder and teats with potable water and dry with a single service paper towel.
- Appropriately clean teat dipper after each milking.

When cow is being dried off, the teats should be dipped once a day at least 3 or 4 days after the last milking.

When freshening cows, begin dipping teats with Double Dip twice a day for 3-4 days before calving.

DIP SOLUTION SHOULD BE CHANGED IF IT BECOMES VISIBLY DIRTY, IF SEDIMENT IS PRESENT, OR IF THE AMBER COLOR FADES NOTICEABLY.

Double Dip may aid in controlling the spread of organisms that cause mastitis.

Do not use for cleaning or sanitizing milking equipment.

Distributed by: EASTERN CROWN, INC.
CHEMICAL DIVISION
PO BOX 850, VERNON, NY 13476
877.829.3505

CAUTION!
Can be severely irritating to eyes. Can be irritating to skin and harmful if swallowed.

First Aid
ORAL INGESTION: If swallowed, dilute by giving several glasses of water or milk. Do not give anything by mouth to an unconscious person. Do not induce vomiting. If vomiting does occur, repeat giving several glasses of water or milk and consult a physician.

EYE CONTACT: Flush eyes with plenty of clean running water for at least 15 minutes. If irritation occurs and persists, get medical attention.

SKIN CONTACT: Wash thoroughly. If irritation occurs and persists, get medical attention.

INHALATION: Remove victim to fresh air.

STORAGE AND DISPOSAL:
DO NOT ALLOW TO FREEZE.
Store in cool dry area. Shake well before using. Keep container tightly closed when not in use. Do not reuse empty container. Rinse container before disposing in a safe manner. Avoid storing near products containing chlorine.
NFEPA=100

NET CONTENT: Gallons

LOT NO.:

EXPIRATION DATE:

ECI

DOUBLE DIP
IODINE TEAT DIP
NDC 060357-1010

ACTIVE INGREDIENT: Iodine (titratable): 1.0%
OTHER INGREDIENTS: 99.0% Contains Glycerin and Lanolin.
Contains No Nonylphenoxy Poly Ethanol (NPE)

KEEP OUT OF REACH OF CHILDREN.
CAUTION
Harmful if Swallowed • Eye Irritant
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS.

Directions for Use:
Use at full strength. Do not dilute.

Before starting or continuing to use teat dip, consult your veterinarian if cow's teats are sore or chapped. ECI Double Dip is manufactured for double use as both a before and after milking dip.

Before Milking:

- Remove excess organic debris from teats and forestrip. If water is used, use minimal amount. Thoroughly dry with a single service towel.
- Check foremilk for abnormalities.
- Dip or spray entire cow's teats with ECI DOUBLE DIP.
- Allow 15 to 30 seconds contact time.
- Dry thoroughly, with clean, single service paper towel.
- Attach milking machine.

After Milking:

- Immediately after milking, dip or spray each entire teat with ECI DOUBLE DIP. Allow to dry. **Do not wipe.** Before next milking, wash cow's udder and teats with potable water and dry with a single service paper towel.
- Appropriately clean teat dipper after each milking.

When cow is being dried off, the teats should be dipped once a day at least 3 or 4 days after the last milking.

When freshening cows, begin dipping teats with Double Dip twice a day for 3-4 days before calving.

DIP SOLUTION SHOULD BE CHANGED IF IT BECOMES VISIBLY DIRTY, IF SEDIMENT IS PRESENT, OR IF THE AMBER COLOR FADES NOTICEABLY.

Double Dip may aid in controlling the spread of organisms that cause mastitis.

Do not use for cleaning or sanitizing milking equipment.

Distributed by: EASTERN CROWN, INC.
CHEMICAL DIVISION
PO BOX 850, VERNON, NY 13476
877.829.3505

DOUBLE DIP

iodine teat dip solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:60357-1010
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	10 g in 1 L

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
LANOLIN (UNII: 7EV65EAW6H)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	brown (liquid)	Score	
Shape		Size	

Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60357-1010-1	3.79 L in 1 BOTTLE, PLASTIC		
2	NDC:60357-1010-2	18.93 L in 1 PAIL		
3	NDC:60357-1010-3	56.78 L in 1 DRUM		
4	NDC:60357-1010-4	208.20 L in 1 DRUM		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
unapproved drug other		10/21/2004		

Labeler - Eastern Crown Inc. (013337118)

Establishment			
Name	Address	ID/FEI	Business Operations
Thatcher Company of New York		043921993	api manufacture, manufacture

Revised: 9/2024

Eastern Crown Inc.