DOUBLE DIP- iodine teat dip solution Eastern Crown Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

ECI Double Dip Iodine Teat Dip

DIRECTIONS FOR USE

Use at full strength. Do not dilute.

Before starting or continuing to use teat dip, consult your veterinarian if cow's teats are sore or chapped.

ECI Double Dip is manufactured for double use as both a before and after milking dip.

Before Milking:

- Remove excess organic debris from teats and forestrip. If water is used, use minimal amount. Thoroughly dry with a single service towel.
- Check foremilk for abnormalities.
- Dip or spray entire cow's teats with ECI DOUBLE DIP.
- Allow 15 to 30 seconds contact time.
- Dry thoroughly, with clean, single service paper towel.
- Attach milking machine

After Milking:

- Immediately after milking, dip or spray each entire teat with ECI DOUBLE DIP. Allow to dry. **Do not wipe**. Before next milking, wash cow's udder and teats with potable water and dry with a single service paper towel.
- Appropriately clean teat dipper after each milking.

When cow is being dried off, the teats should be dipped once a day at least 3 or 4 days after the last milking.

When freshening cows, begin dipping teats with Double Dip twice a day for 3-4 days before calving.

DIP SOLUTION SHOULD BE CHANGED IF IT BECOMES VISIBLY DIRTY, IF SEDIMENT IS PRESENT, OR IF THE AMBER COLOR FADES NOTICEABLY.

Double Dip may aid in controlling the spread of organisms that cause mastitis.

Do not use for cleaning or sanitizing milking equipment.

STORAGE AND DISPOSAL:

DO NOT ALLOW TO FREEZE.

Store in cool dry area. Shake well before using. Keep container tightly closed when not in use. Do not reuse empty container. Rinse container before disposing in a safe manner. Avoid storing near products coontaining chlorine.

CAUTION!

Can be severely irritating to eyes. Can be irritating to skin and harmful if swallowed.


First Aid

ORAL INGESTION: If swallowed, dilute by giving several glasses of water or milk. Do not give anything by mouth to an unconscious person. Do not induce vomiting. If vomiting does occur, repeat giving several glasses of water or milk and consult a physician.

EYE CONTACT: Flush eyes with plenty of clean running water for at least 15 minutes. If irritation occurs and persists, get medical attention.

SKIN CONTACT: Wash thoroughly. If irritation occurs and persists, get medical attention.

INHALATION: Remove victim to fresh air.



DOUBLE DIP

iodine teat dip solution

Product Information

Product Type		OTC ANIMAL DRUG	item Co	de (Source)	N	IDC:60357-1010	
Route of Administra	ation	TOPICAL					
Active Ingredient	/Active	Moiety					
Ingredient Name				Basis of S	Streng	th Strengt	
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)				IODINE		10 g in 1 L	
Inactive Ingredie	nts						
Ingredient Name						Strength	
GLYCERIN (UNII: PDC6A	3C0OX)						
LANOLIN (UNII: 7EV65E	AW6H)						
WATER (UNII: 059QF0KO0R)							
Product Characte	eristics						
Color brown		wn (liquid) Score					
Shape			Size				
Flavor			Imprint Code				
Contains							
Dackaging							
Packaging	Deelse		Maulsatius		Mari	esting Fuel Det	
# Item Code			Marketing	g Start Date	Mark	ceting End Dat	
1 NDC:60357-1010-1		L BOTTLE, PLASTIC					
2 NDC:60357-1010-2	18.93 L in						
3 NDC:60357-1010-34 NDC:60357-1010-4	56.78 L in 208.20 L i						
+ NDC:00557-1010-4	200.20 L I						
Marketing Inf	ormat	ion					
Marketing	Applica	lication Number or Monograph Citation		Marketing Start Date		Marketing End Date	
Category							

Labeler - Eastern Crown Inc. (013337118)

Establishment								
Name	Address	ID/FEI	Business Operations					
Thatcher Company of New York		043921993	api manufacture					

Eastern Crown Inc.