

TYLENOL EXTRA STRENGTH- acetaminophen tablet
R J General Corporation

TYLENOL® EXTRA STRENGTH

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

if a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"> ▪ take 2 caplets every 6 hours while symptoms last ▪ do not take more than 6 caplets in 24 hours, unless directed by a doctor ▪ do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store between 20-25°C (68-77°F)
- **do not use if pouch is torn or damaged**

Inactive ingredients

carnauba wax ¹, corn starch ¹, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, modified starch ¹, polyethylene glycol ¹, powdered cellulose, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, titanium dioxide

1 contains one or more of these ingredients

Questions or comments?

call **1-877-895-3665**(toll-free) or **215-273-8755**(collect)

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA

Repackaged and distributed by:

RJ General
2024 Northwest Drive
Cincinnati OH 45231

PRINCIPAL DISPLAY PANEL - 500 mg Caplet Pouch Box

TYLENOL[®]
FOR ADULTS

Acetaminophen
Pain Reliever
Fever Reducer

Extra Strength

50 Pouches of
2 Caplets each
500 mg each

Do not use if pouch is torn or damaged.

TYLENOL[®]

FOR ADULTS

Acetaminophen Pain Reliever
Fever Reducer

Extra Strength



50 Pouches of
2 Caplets each
500 mg each

Do not use if pouch is torn or damaged.

50 Pouches of 2 Caplets each - 500 mg each

TYLENOL[®]

FOR ADULTS



Acetaminophen Pain Reliever
Fever Reducer

Extra Strength



50 Pouches of
2 Caplets each
500 mg each

Contains No Aspirin




How can we help?
 MSG & DATA RATES MAY APPLY
 TEXT HELP FOR HELP
 TEXT STOP FOR STOP
 SNAP THIS CODE
 OR TEXT TYLENOL
 TO 87719

1-877-895-3665



**Important: Read all product information before using.
Keep this package for important information.**

Drug Facts

Active ingredient (in each caplet) Purpose
 Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - backache
 - toothache
 - premenstrual and menstrual cramps
 - headache
 - minor pain of arthritis
 - muscular aches
 - temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash
 if a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over	take 2 caplets every 6 hours while symptoms last do not take more than 6 caplets in 24 hours, unless directed by a doctor do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store between 20-25°C (68-77°F)
- do not use if pouch is torn or damaged

Inactive ingredients


carnauba wax*, corn starch*, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, modified starch*, polyethylene glycol*, powdered cellulose, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, titanium dioxide
 *contains one or more of these ingredients

Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

TYLENOL[®]
 FOR ADULTS
 Acetaminophen Pain Reliever
 Fever Reducer
Extra Strength





TYLENOL EXTRA STRENGTH

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70264-027(NDC:67414-449)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
--------------	-------	--------------	----------

Shape	OVAL	Size	18mm	
Flavor		Imprint Code	TYLENOL500	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70264-027-01	50 in 1 BOX	05/01/2021	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	05/01/2021		

Labeler - R J General Corporation (122542830)

Establishment			
Name	Address	ID/FEI	Business Operations
R J General Corporation		122542830	repack(70264-027)

Establishment			
Name	Address	ID/FEI	Business Operations
Johnson & Johnson		878046358	manufacture(70264-027)

Revised: 11/2025

R J General Corporation