

**EXTRA STRENGTH PAIN RELIEF- acetaminophen suspension  
GERI-CARE PHARMACEUTICAL CORP**

---

**GerCare- APAP SUS Extra 206**

**Active ingredient (in each 15mL Tablespoonful)**

Acetaminophen 500 mg

**Purpose**

Pain reliever/Fever reducer

**Uses**

- temporarily relieves minor aches and pains
- temporarily reduces fever

**Warnings**

**Liver Warning:** This product contains acetaminophen. The maximum daily dose of this product is 6 tablespoonfuls in 24 hours.

Severe liver damage may occur if you take

- more than 8 tablespoonfuls (4,000 mg of acetaminophen) in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product: Do not exceed recommended dose.**

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

**If pregnant or breastfeeding,** ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults/children even if you do not notice any signs or symptoms.

### **Directions**

- do not take more than directed

Adults and children 12 years of age and older	Take 1 to 2 Tablespoons every 4 to 6 hours as needed; not more than 6 Tablespoons in 24 hours. Do not take for more than 10 days unless directed by a doctor
Children under 12 years	ask a doctor

### **Other information**

- store at room temperature 15°- 30°C (59°-86°F)
- protect from freezing
- protect from light
- **TAMPER-EVIDENT:** Do not use if foil seal over bottle opening is torn, broken or missing.

### **Inactive ingredients**

artificial and natural cherry flavor, citric acid, FD&C Red #40, methylparaben, polyethylene glycol, propylene glycol, propylparaben, purified water, sodium citrate, sucralose


### **Questions or comments?**

**1-800-540-3765**

### **PRINCIPAL DISPLAY PANEL**

<b>Drug Facts</b>	
<b>Active ingredient</b> (in each 15mL Tablespoonful)	<b>Purposes</b>
Acetaminophen 500 mg.....	Pain Reliever / Fever Reducer
<b>Uses</b>	
• temporarily relieves minor aches and pains • temporarily reduces fever	
<b>Warnings</b>	
<p><b>Liver Warning:</b> This product contains acetaminophen. The maximum daily dose of this product is 6 tablespoonfuls in 24 hours. Severe liver damage may occur if you take • more than 8 tablespoonfuls (4,000 mg of acetaminophen) in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product</p> <p><b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.</p>	
<p><b>Do not use</b> • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</p> <ul style="list-style-type: none"> <li>• if you are allergic to acetaminophen or any of the inactive ingredients in this product</li> </ul> <p><b>Ask a doctor before use if you have liver disease.</b></p> <p><b>Ask a doctor before use if you are taking the blood thinning drug warfarin.</b></p> <p><b>When using this product: Do not exceed recommended dose.</b></p> <p><b>Stop use and ask a doctor if</b> • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present. These could be signs of a serious condition.</p>	

NDC 57896-206-08

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults/children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed
- adults and children 12 years of age and older take 1 to 2 tablespoonfuls every 4 to 6 hours as needed; not more than 6 tablespoonfuls in 24 hours. Do not take for more than 10 days unless directed by a doctor.
- Children under 12 years: ask a doctor

**Other information**

- store at 20°C-25°C (68°F-77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- protect from freezing • protect from light
- **TAMPER-EVIDENT:** Do not use if foil seal over bottle opening is torn, broken or missing.

**Inactive ingredients:** artificial and natural cherry flavor, citric acid, FD&C Red #40, methylparaben, polyethylene glycol, propylene glycol, propylparaben, purified water, sodium citrate, sucralose

**Questions or comments?** 1-800-540-3765

\*This product is not manufactured or distributed by the owner of the registered trademark Tylenol®. Code # L-23  
Dist. By: Geri-Care Pharmaceuticals Corp. Rev.:11/25  
1295 Towbin Avenue  
Lakewood, NJ 08701 3 57896 20208 8  
Lot#  
Exp.  
Date:

**EXTRA STRENGTH PAIN RELIEF**

acetaminophen suspension

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:57896-206
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg in 15 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
METHYLPARABEN (UNII: A2I8C7H9T)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	

**Product Characteristics**

<b>Color</b>	red	<b>Score</b>
--------------	-----	--------------

<b>Shape</b>	<b>Size</b>
<b>Flavor</b>	<b>Imprint Code</b>
<b>Contains</b>	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-206-08	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2019	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/01/2019	

## Labeler - GERI-CARE PHARMACEUTICAL CORP (611196254)

Revised: 12/2025

GERI-CARE PHARMACEUTICAL CORP